

# Samaritan Pacific Communities Hospital

## 2016 Community Health Needs Assessment



**Project Team** (alphabetical):

Peter Banwarth, MS  
Valerie Barnhill, MPH Intern  
Miyuki Blatt, MPH Intern  
Ann Brown, MBA  
Savannah Carrico, MPH intern  
Andy Chuinard, MPH  
Jessica Deas, MPH  
Gerald Dyer, MPH  
Charlie Fautin, MPH, RN  
Brandan Kearny, Consultant  
Ruby Kiker, MPH  
JoAnn Miller, MS  
Megan Patton-Lopez, PhD, RD  
Whitney Schumacher, MPH intern  
Lauren Zimbelman, MPH

**For Further Information**

Samaritan Health Services  
Community Health Promotion  
JoAnn Miller, Director  
815 NW 9<sup>th</sup> Street  
Corvallis, OR 97330  
541-768-7330  
jomiller@samhealth.org

Regional Health Assessment  
530 NW 27<sup>th</sup> St.  
Corvallis, OR 97339  
LBLRHA@co.benton.or.us  
541-766-3547  
Fax: 541-766-6142

**Table of Contents**

**Chapter 1. Introduction and Overview ..... 1**

- Samaritan Pacific Communities Hospital ..... 1
- How to Use This Document ..... 4

**Chapter 2. People ..... 6**

- Health Disparities ..... 6
- Population Overview ..... 7
- Demographics: Population by Age and Sex ..... 7
- Growing Diversity ..... 8
- Disabilities ..... 11
- Aging Population ..... 11
- Conclusion ..... 11

**Chapter 3. Environment ..... 13**

- Natural Environment ..... 13
- Recreation and outdoor spaces ..... 14
- Fluoridated Water ..... 15
- Human-made Environment ..... 15
- Tobacco-free Spaces ..... 16
- Transportation ..... 16
- Conclusion ..... 18

**Chapter 4. Social Determinants of Health ..... 19**

- Income, Poverty, and Economic Challenges ..... 19
- Education ..... 23
- Food Security ..... 25
- Housing and Home Ownership ..... 27
- Conclusion ..... 29

**Chapter 5. Access to Medical Care ..... 30**

- Health Insurance Coverage ..... 30
- Cost of Medical Care ..... 32
- Access Capacity ..... 33
- Safety Net Services & Community Benefits ..... 33
- Medical Services ..... 34
- Conclusion ..... 41

<b>Chapter 6. Morbidity and Mortality .....</b>	<b>42</b>
Leading Causes of Death in Lincoln County .....	42
Chronic Disease and Conditions .....	43
Heart Disease and Stroke.....	46
Diabetes .....	47
Alzheimer’s Disease .....	47
Arthritis .....	47
Asthma .....	48
Infectious Diseases.....	48
Injury and Violence .....	51
Mental Health Conditions.....	52
Conclusion.....	53
<b>Chapter 7. Health Across the Life Course.....</b>	<b>54</b>
Maternal and Infant Health .....	55
Childhood and Adolescence.....	62
Oral Health .....	68
Alcohol, Tobacco, and Prescription and Illicit Drug Abuse .....	69
Adults .....	71
Older Adult Health .....	77
Conclusion.....	78
<b>Conclusion. Meeting Challenges with Resources/Evaluation Impact .....</b>	<b>80</b>

# Chapter 1

## Introduction and Overview

### Samaritan Pacific Communities Hospital

Samaritan Pacific Community Hospital, doing business as Samaritan Pacific Health Services, was established in 1952 as a 17-bed acute care facility. On July 1, 2003 SPCH received its designation as a 25-bed Critical Access Hospital. SPCH serves all of Central Lincoln County and is part of the Pacific Communities Health District. The hospital recently opened the first dialysis center in the Newport area. Other services include emergency care, obstetrics, critical care, diabetes education, health and medical/surgical services.

Samaritan Pacific Communities Hospital operates within the Pacific Communities Health District. The mission of the Pacific Communities Health District is to deliver high quality health care services to the residents and visitors of the District and its surrounding communities. A wide range of health care services comprises the Samaritan Pacific Communities Hospital organization, which delivers continuous service to a very diverse population. The service area stretches from Depoe Bay to mid-Lincoln Beach, Yachats and east to Eddyville.

As a non-profit hospital, we are committed to our motto “building healthier communities together.”

The 2016 Samaritan Pacific Communities Hospital Community Health Needs Assessment (CHNA) is the result of many dedicated hours of research, working in collaboration with community partners and agencies, leaders, and local residents in Central Lincoln County.

The World Health Organization defines health as a “state of complete physical, mental and social well-being and not merely the absence of disease or infirmity.”<sup>1</sup> Health is not just about individuals, but also includes families, communities, and systems, and is a result of the interaction of complex networks of conditions and factors. Health starts long before illness occurs and is impacted by where and how we live, learn, work, play, worship and age.

This CHNA incorporates this definition of health by describing a wide array of information about the conditions and factors affecting people’s health across Central Lincoln County as well as indicators of health status.

### Assessment Goals and Objectives

The Samaritan Pacific Communities Hospital Community Health Needs Assessment (CHNA):

- Identifies and gathers health status indicators in order to determine the current health status of the Central Lincoln community;

- describes areas for potential future health improvement while building upon ongoing community knowledge and efforts;
- recognizes and highlights the need for more detailed local data; and
- is a collaborative process that incorporates a broad range of community voices.

CHNA data informs Samaritan Pacific Communities Hospital’s organizational decision-making, the Samaritan Pacific Communities Hospital’s Community Benefit Implementation Plan, and other community and hospital efforts.

## Needs Assessment Organization

The Community Health Needs Assessment is presented in eight chapters. An appendix of key informant interviews, focus groups, and survey results is available as a separate document.

Chapter 1: *Introduction and Overview*, including methodology and limitations.

Chapter 2: *People*, describing the people of Central Lincoln County, including population demographics as well as a look at how the community has changed over time.

Chapter 3: *Environment*, which includes information about the physical spaces in which we live, work, and play.

Chapter 4: *Social Determinants of Health*, which includes the social, economic, and community factors that influence health.

Chapter 5: *Access to Medical Care*, exploring how we define and measure the ability of those in our community to get the medical care they need.

Chapter 6: *Morbidity and Mortality*, which covers a number of related health outcomes, from chronic conditions to violence and injury.

Chapter 7: *Health Across the Life Course*, exploring the ways in which individuals and communities act to protect and improve health at different stages in life.

Chapter 8: *Conclusion: Meeting Challenges with Resources/Evaluation Impact*, discussing how this data can be used to understand the health of Central Lincoln County and recognize opportunities for positive changes to improve the health of the entire community.

Appendix (separate): *Community Perceptions on the Health of Central Lincoln County*, reporting on the results of key informant interviews, focus groups, and a community survey on perceptions of the health of the community.

## Methodology

The CHNA is comprised of secondary and primary data. Data from secondary sources were identified through meeting with community partners, and through preexisting publications (e.g.

community health assessments). In addition, data sources were identified through literature research to include data from local state and national levels. A variety of community partners were involved throughout this process. Staff conducted both in-person and phone presentations and consultations with members of regional and county-level governmental, nonprofit, and health system organizations. In addition, members of state and local research communities were contacted.

This process has included:

- Engaging community stakeholders and partners in the process of issue identification, data collection, data interpretation, editing, and dissemination of results;
- obtaining updated secondary data for the Central Lincoln County;
- synthesizing existing data reports; identifying areas in which more information is needed, and including data from other sources which address these gaps;
- identifying health needs and assets that will inform additional local planning processes, including county-level Community Health Improvement Plans, Public Health Division strategic planning, public health accreditation, and health care transformation initiatives, among others; and
- consulting state and national resources for guidance in the development of this community health assessment, including the following: Oregon Health Authority technical reports (e.g. health equity,<sup>2</sup> asthma,<sup>3</sup> chronic disease prevention<sup>4</sup>); the Centers for Disease Control and Prevention's data set directory of social determinants of health at the local level;<sup>5</sup> King County's Equity and Social Justice Annual Report;<sup>6</sup> and the Statewide Health Assessment of Minnesota.<sup>7</sup>

## Limitations

While the CHNA identifies many critical issues pertaining to community health, it is not inclusive of all health-related issues. As a result, it should not be considered a formal study or research document investigating the causes of each issue raised or providing a detailed analysis of the data.

When considering the many factors that contribute to health, data are lacking in part because respective theoretical models are still being developed. In addition, conclusions, hypotheses, and interpretations of the interactions between the many factors that contribute to health may not be included, in part because the underlying structures of these interactions are still not fully understood.

## Gaps in Data

Recognizing and highlighting the need for more detailed local data was a key objective of this assessment. As mentioned throughout the document, data for the Central Lincoln County were often not available for particular demographics, such as age, income, education-level,



race/ethnicity, or zip code. This greatly limited the ability to explore differences or disparities within particular sub-populations.

When race/ethnicity data are gathered, analysis may be further limited due to a lack of data stratification by more specific racial categories, such as U.S.-born versus foreign-born for the Latino population, or the many ethnicities and cultures represented in the category of Asian-Pacific Islander. There are limited data on disparate populations in the community; however, as highlighted in later chapters, their needs and barriers to health and health care are likely to be greater than those of the population at large.

Throughout the document, Oregon state-wide data are provided to illustrate trends, especially among vulnerable populations, when county level data are not available. It is important to note, however, that national or state-wide rates, trends, and patterns may not necessarily reflect the reality of particular communities or counties. As community partners continue to gather information to inform their practices and services, it is important to collect demographic data (i.e. zip codes, level of education, etc.) so that more accurate information can be used to inform future health improvement planning and other public health initiatives.

## **How to Use This Document**

### **Central Lincoln County versus Lincoln County**

Samaritan Pacific Communities Hospital serves the people of Central Lincoln County, comprising the communities of Newport, Siletz, Toledo, Depoe Bay, Waldport, and Yachats, as well as unincorporated and rural areas of Central Lincoln County. Demographic and economic data was largely available for Central Lincoln County; however, most health status data is not reported at a sub-county level. When data is specific to central Lincoln County, it is identified as such. When data is reported for all Lincoln County, this is noted as well. All of Lincoln County includes the communities of Lincoln City, Lincoln Beach, Gleneden Beach, and Otis.

### **Timeframes for Data**

This report attempts to balance the importance of comparing data from common years with the goal of presenting the most recent data. Different data sources update and release data on independent timeframes. The U.S. Census Bureau is the main source of data for demographic and socioeconomic information used in this report. The most precise data available for county-level demographic and socioeconomic data is the Census Bureau's American Community Survey (ACS) 2011-2013 three-year aggregates. This aggregation combines data from the three years in order to produce more accurate estimates. The most precise data available for sub-county estimates (the South Lincoln region), is a five-year aggregate from 2010-2014.

In an effort to compare data from common years, many statistics reported are from 2013, even if more recent data is available. These statistics reflect measures of health that have historically



changed gradually, so differences between 2013 and the present are likely to be minor. However, some measures of health have changed greatly in the past two years, such as the implementation of the Affordable Care Act in 2014, which had huge impacts on insurance coverage rates and Medicaid membership. In this case and for other rapidly changing measures, more contemporary data is reported in order to best reflect current health status and the current health system.

As with the ACS 2011-2013 3-year aggregates, many data sources aggregate statistics over a number of years to improve the reliability of the estimates. A common example of this is reporting the incidence (number of new cases) of cancer. For example, in the state of Oregon there were approximately 98,860 new cases of cancer in Oregon between 2008 and 2012. This statistic is reported as an incidence of 448 cases for every 100,000 people. This means that *each year*, for every 100,000 people in Oregon there were 448 cancer diagnoses. It does not mean that 448 cases per 100,000 people were diagnosed over the course of 5 years.

# Chapter 2

## People

The history of Central Lincoln County begins with the Native American tribes that have lived in the region for thousands of years. Native Americans lived in the valleys and the hills, along rivers and oceans.

Contact with non-native groups began with trappers and explorers in the late 18<sup>th</sup> century, then with pioneers and settlers who moved to the Oregon Territory during the mid-1800s. In 1855, the United States established a 1.3 million acre reservation in what is now Lincoln County. The U.S. government moved many of the coastal and Willamette Valley tribes to this reservation, which at the time included Yaquina Bay. As a result, Lincoln County was characterized by a large Native American presence and interaction between Native Americans and immigrants.

Over the next 150 years, Lincoln County incorporated, grew in population, and developed strong local industries. Today, natural resources are highly valued in Lincoln County. Industries such as commercial fisheries, logging, tourism, and agriculture are drivers of the local economy. The county is also home to state and national research centers, and the Confederated Tribe of the Siletz Indians.

### Health Disparities

In the discussion of residents of Central Lincoln County, it is important to recognize that specific subpopulations may experience worse health outcomes than the general population. These include: individuals with mental health issues; homeless; rural residents; racial or ethnic groups; those with low educational attainment; persons living in poverty; and lesbian, gay, bisexual, transgender, or queer (LGBTQ) persons, among other subpopulations.

For example, across the United States, rates of illness for adults in their 30s and 40s with lower incomes and education levels are comparable to affluent adults in their 60s and 70s. College graduates can expect to live at least five years longer than those who have not finished high school, and almost two years longer than those who did not finish college.<sup>8</sup> People with mental health conditions and/or substance abuse issues in Oregon lose an average of 34.5 potential life years.<sup>9</sup> Understanding health disparities in the context of Lincoln County is critical to improving health across the county.

This chapter on the people of Central Lincoln County serves as the foundation for understanding the different characteristics of the people in the county. Permanence of residency has important effects on health, but at this time there is not data detailing the mobility of residents within the county and to or from other areas. This makes it more difficult to provide an accurate picture of changing population dynamics. However, thanks in part to

comprehensive data collected by the U.S. Census Bureau, we have a strong understanding of the residents of Central Lincoln County. This chapter will detail Central Lincoln County population, its population distribution, population centers, and population characteristics of age, gender, race, ethnicity, disability and diversity.

## Population Overview

Central Lincoln County is home to 31,668 residents.<sup>10</sup> Most of these residents live in Newport (pop. 10,095), Toledo (pop. 3,485) and Waldport (pop. 2,060), with other residents in smaller towns and rural communities.

In 2013, there were 14,085 households in Central Lincoln County. Household distribution follows roughly the same pattern as overall population distribution across the county. The average household size in Central Lincoln County was 2.2 people. Families made up 62 percent of the households. This figure includes both married couple families and other family households.<sup>11</sup> Non-family households made up 38 percent of all homes in Central Lincoln County. Most non-family households are composed of people living alone, but some are people living in households in which no one is related to the head of household.<sup>12</sup>

In Central Lincoln County, 20 percent of all households have one or more people under the age of 18. Thirty-eight percent of all Lincoln County households have one or more people 65 years and over.<sup>13</sup>


## Veterans

The 2009-2013 American Community Survey (ACS) report the veteran population in central Lincoln County at 4,003.<sup>14</sup> Veterans are defined as men and women who have previously served on active duty in the U.S. Army, Navy, Air Force, Marine Corps, Coast Guard, or who served in the U.S. Merchant Marine during World War II.<sup>15</sup> This amounts to approximately 15 percent of Central Lincoln County's civilian population ages 18 years and older. As this population ages, the number of individuals with veteran status is expected to decrease over time.

## Demographics: Population by Age and Sex

Based on U.S. 2013 Census data, the percentage of males and females in Central Lincoln County is approximately equal in most age groups.<sup>16</sup> Within Central Lincoln County, children under 18 years of age constitute 17 percent of the population and the age group of adults 65 years and older constitutes 23 percent of the population. The median age in Central Lincoln County is ranges from 35 to 65 years, depending on the Census tract.<sup>17</sup> From 2000 to 2014, the population of Central Lincoln County grew 4.5 percent.<sup>18</sup>

Figure 2.1: Central Lincoln County population by age group, 2014



Source: U.S. Census Bureau, American Community Survey 5-year estimates, 2010-2014

## Growing Diversity

### Native and Foreign Born

In 2013, 94.2 percent of the people living in Central Lincoln County were native residents of the United States.

### Race/Ethnicity

In an increasingly global view of health and understanding of research outlining the social constructs of race and ethnicity, a culturally sensitive definition of race should be considered. The inclusion of individuals to self-identify as two or more races has been adopted almost universally across other agencies collecting and reporting demographic data. It is important to understand the data for individuals along the lines of racial divide as later issues of health disparities will be presented. Without understanding the populations impacted by these health disparities, health authorities would be limited in their ability to address the specific issues creating the disparities.

U.S. Office of Management and Budget defines race and ethnicity categories accordingly:

**White** – people having origins in Europe, the Middle East, or North Africa.

**Black or African-American** – people having origins in the black racial groups in Africa.

**Asian** – people having origins in any of the original peoples of the Far East, Southeast Asia or the Indian subcontinent.

**Native Hawaiian or Other Pacific Islander** – people having origins in any of the original people of Hawaii, Guam, Samoa, or other Pacific Islands.

**American Indian or Alaska Native** – people having origins in any of the original peoples of North or South America (including Central America), and who maintain a tribal affiliation or community attachment.

**Multiracial** – people having origins in two or more of the federally designated racial categories.

**Hispanic or Latino** – A person of Cuban, Mexican, Puerto Rican, South or Central America, or other Spanish culture or origin, regardless of race.<sup>19</sup>

In this report, the non-Hispanic categories are used for races. For example, the category denoted White includes white, non-Hispanic individuals.

As shown in Table 2.1, the largest ethnic population in Central Lincoln County is Latinos (7.8 percent).<sup>20</sup> The Latino population increased by three percent from 2000 to 2011.<sup>21,22,23</sup> In Central Lincoln County, the predominant non-white racial group are individuals who identify as being multi-racial (4.7%). In general, Central Lincoln County’s population is less diverse than that of the state of Oregon (Figure 2.2).


Table 2.1: Population by Race/Ethnicity Central Lincoln County 2011-2013\*

Race and Ethnicity	Central Lincoln County	
	Total Population	Percent
White	27,573	87%
Hispanic or Latino	2,479	7.8%
Asian	193	0.6%
Two or more races	1,499	4.7%
American Indian and Alaska Native	1,032	3.2%
Black or African American	135	0.4%
Hawaiian and Pacific Islander	61	0.2%
Other race	1,175	3.7%

Source: U.S. Census Bureau, American Community Survey 3-year estimates, 2011-2013, Table DP05

\*Columns may not sum to 100% due to rounding error

Figure 2.2: Population by Race/Ethnicity in Central Lincoln County, 2011-2013



Source: U.S. Census Bureau, American Community Survey 5-year estimates, 2010-2014

## K-12 Population

Lincoln County encompasses a single school district with 17 public schools serving 5,237 students during the 2013-2014 school year. This represents an increase in the number of students by 1.1 percent since 2010.<sup>24</sup> These statistics do not include private schools or homeschooled students. In Table 2.2 below, the racial and ethnic diversity for the county school district is compared to the region overall.

Table 2.2: Lincoln County School District by race/ethnicity, 2013-2014

County	White	Hispanic/ Latino	Asian Pacific Islander	Multi- Ethnic	American Indian/ Alaskan Native	African American	Total Minority
<b>Lincoln County SD</b>	66.4%	18.5%	8.6%	6.9%	6.3%	< 1.0%	33.7%

Source: Oregon Department of Education, Student Ethnicity statistics, academic year 2013-2014

## Language Spoken at Home

2011-2013 U.S. Census data reports that 2.5 percent of Central Lincoln County residents who are at least 5 years old spoke English less than very well.

## Disabilities

Understanding and measuring disability is a very complex task. The complexity comes from the fact that the definition of “disability” includes a number of populations, and because the definition is still being discussed and further developed. Definitions of disabilities from a source such as the World Health Organization (WHO) can help shed light on the particular health issues facing these populations, but it must be noted that this definition is not the same as that used to gather many types of data.

Disability itself is not an indicator of poor health—rather, disability can (and often does) become a barrier to employment, adequate housing, social inclusion, transportation, access to health care, and other essential components of a healthy life.

Mental illness, that substantially limits one or more major life activities, is also included in definitions of disability.<sup>25</sup> This is particularly worth noting, as institutionalized populations generally experience a greater prevalence and severity of mental illness than the broader population. However, these populations are not captured in much of the data collected around disability.<sup>26</sup>

From 2011 to 2013, among the civilian non-institutionalized population, approximately 22 percent reported a disability in Central Lincoln County,<sup>27</sup> where *disability* is defined as a person’s risk of participation limitation when he or she has a functional limitation or impairment.<sup>28</sup> Disability encompasses many different conditions; for instance, the most common disability in all of Lincoln County among those aged 5-64 is cognitive difficulty, with ambulatory difficulty ranking the highest for the 65 and older population. The prevalence of disability increases with age, from 1 percent of people under 5 years of age in Lincoln County, up to 14 percent for 18 to 64 years of age, and 41 percent of those 65 and over.

## Aging Population

Among those living in Central Lincoln County, 23 percent are 65 years of age and over, compared with 14 percent in Oregon overall.<sup>29</sup> A number of health issues, needs, and concerns are associated with an aging population.

## Conclusion

In order to understand the health of Central Lincoln County, it is vital to understand the people who live here. Differences in age, race or ethnicity, and geography all influence health. Vulnerable populations, such as individuals with disabilities or older adults, merit further description, both because they may require different services, and also because they may present different health concerns. The people of Central Lincoln County are growing more diverse and represent many different groups, such as students, Native Americans, and retirees. The history of Central Lincoln County has shaped the residents of this county into its makeup



today. In exploring the many determinants of health, it is evident that the people of Central Lincoln County are deeply connected with the environments in which they live. The next chapter explores these environments and the effects they have on the health of Central Lincoln County.

# Chapter 3

## Environment

Human beings interact with their environment in everything they do. Some of these interactions have the potential to improve health, while others can negatively impact it. The natural environment is made up of the interactions of air, water, open spaces, and weather or geologic activity. The human-made environment consists of homes, communities, and infrastructure. These two environments are closely linked in their effects on human health. Humans benefit from clean water and air, places to exercise and enjoy the outdoors, safe living and working spaces, and opportunities to engage in healthy behaviors such as active commuting and consuming healthy food. However, when an environment lacks these characteristics, the complex interactions of health and environment can worsen health issues. Poor air quality can raise the risk of asthma, heart attack, or stroke;<sup>30</sup> poorly designed communities can limit opportunities for recreation or access to quality food;<sup>31</sup> and infrequent but intense natural disasters can disproportionately affect vulnerable populations.

Central Lincoln County has many strengths when it comes to environmental factors and health opportunities. Residents of Central Lincoln County value open spaces for recreation, clean air, and clean water. At the same time Central Lincoln County faces challenges with food access and transportation issues. An understanding of the natural and human-made environments forms a foundation for an analysis of the health of Central Lincoln County. Most of the environmental data presented in this chapter is at the county geographic level.

### Natural Environment

The natural environment changes slowly and usually influences health through long-term, cumulative effects. As a result, many of the data described in this section use longer time frames than elsewhere in this report. Furthermore, unlike many other determinants of health, it can be very difficult or impossible for individuals or health professionals to influence the natural environment on a local scale. Examples include global climate change or natural disasters. What can be controlled are the systems and practices put in place to react and adapt to the natural environment in order to improve health.

### Terrain and Natural Resources

Lincoln County covers an area of 978 square miles. It is approximately 60 miles north to south and between 20 and 40 miles east to west. The western border of the Lincoln County is the Pacific Ocean, while it is bordered on the east by Benton and Polk Counties, on the north by Tillamook County, and on the south by Lane County. Primary land cover types include mixed Douglas-fir coniferous forests, oceanfront and beaches, agricultural land, and coastal temperate

rainforests. Elevations range from sea level along the coast 3,400 feet in the Coast Mountain range, which forms the eastern half of Lincoln County.

Natural resources are highly valued in Lincoln County; many residents depend on commercial fisheries, logging, tourism, and coastal agriculture. Newport is home to one of the most important fishing ports in the United States. It is also utilized by many local agencies including the U.S. Coast Guard, the National Oceanic and Atmospheric Administration, Oregon Department of Fish and Wildlife, and Oregon State University, among others.

## Annual weather patterns

Lincoln County's location along the Pacific Coast moderates its annual weather patterns. Temperatures typically range from 40 to 80 degrees Fahrenheit over the course of the year.<sup>32</sup> There is an average of 187 days of precipitation per year, which falls as rain in most of the county year round. Lincoln County receives an average of between 70 and 100 inches of rain per year depending on location, with 30 or more heavy rainfall events (one or more inches per day) in some parts of the county.<sup>33,34,35</sup>

## Recreation and outdoor spaces

Lincoln County is favored with a great variety of recreational assets and outdoor spaces. The county's coastline encompasses beaches, headlands, tide pools, and estuaries. The shallow and deep ocean off the coast of Lincoln County is frequented by recreational and commercial fishers. In 1913, the governor of Oregon declared all beaches a public highway, which began a legacy of public access to the oceanfront. In 1967, the state legislature mandated free and uninterrupted use of the beaches along Oregon's 362-mile coastline.<sup>36</sup> Along the coastline, there are a large number of state parks with camping and day-use areas, as well as Beaver Creek State Natural Area south of Newport. Farther inland, the Siuslaw National Forest covers 172,000 acres in Lincoln County and an additional 18,000 acres in Benton County. The forest is networked with hiking trails and hosts deer and elk hunts. Many Lincoln County rivers are open to seasonal fishing, including the Yaquina, Alsea, Siletz, and Salmon.

## Recreational Access

Access to recreational facilities and opportunities demonstrates the intersection of natural and human-made environments. Research demonstrates a strong relationship between access to recreational facilities and physical activity among adults and children. Studies have shown that proximity to places with recreational opportunities is associated with higher physical activity and lower obesity levels.<sup>37</sup> Public recreation areas include parks, schools, public forests and trails, beaches, and waterfronts. As mentioned previously, Lincoln County's rural areas are largely accessible to residents.

Recreational opportunities that include walking and bicycling are efficient, low-cost, and available to most anyone. By walking and bicycling, residents can help develop and maintain livable communities, make neighborhoods safer and friendlier, save on motorized transportation costs, and reduce transportation-related environmental impacts, auto emissions, and noise. They can also create transportation system flexibility by providing alternative mobility options, particularly in combination with transit systems. Furthermore, creating walkable and bikeable communities can lead to healthier lifestyles.<sup>38</sup>

Due to the public nature of Oregon's recreational areas, 55 percent of residents in Lincoln County live within one half mile of a public recreation area, as defined by the CDC.<sup>39</sup> This is the 12<sup>th</sup> highest percentage in the state.<sup>40,41</sup>

## Fluoridated Water

Water fluoridation is the controlled addition of a fluoride compound to a public water supply, intended to prevent tooth decay. Community water fluoridation is an evidence-based practice recommended by the Community Preventive Services Task Force based on strong evidence of effectiveness in reducing dental cavities across populations.<sup>42</sup> It is an effective, affordable, and safe way to protect children from tooth decay and is recognized as one of the 10 greatest public health achievements of the 20<sup>th</sup> century.<sup>43</sup> Water fluoridation complements, but does not replace other efforts to improve oral health. Water fluoridation is a valuable tool in addressing oral health disparities, since everyone who can access public water benefits from it regardless of age, income level, or race or ethnicity. As of 2012, Oregon was ranked very low in the United States (48<sup>th</sup> out of the 50 states) for the percentage of people receiving fluoridated water. About 75 percent of the U.S. population served by community water systems received fluoridated water, while about 23 percent of Oregon's public water supplies are fluoridated.<sup>44</sup> This low state fluoridation rate is a direct consequence of some of Oregon's most densely populated regions lacking fluoridation, including Portland and Eugene.

There are currently no public water systems in Lincoln County that fluoridate water.

## Human-made Environment

Human-made (or built) environments contribute to health in a variety of ways. People need schools, workplaces, and homes that do not expose them to physical or chemical hazards and places to walk and recreate outdoors that are clean, safe, and free of debris. They also need access to quality and affordable food and transportation options, as well as the confidence that their local communities have not been contaminated with human-made pollutants.<sup>45</sup>

## Healthy Homes

Indoor environmental quality, as defined by the Centers for Disease Control and Prevention, is the quality of a building's environment in relation to the health and well-being of those who

occupy the space within it. Key factors that influence a structure's indoor environmental quality include dampness and mold in buildings, building ventilation, construction and renovation, chemicals and odors, indoor temperatures, and relative humidity.<sup>46</sup> Buildings in the Central Lincoln County are often exposed to winter storms with winds in excess of 30 mph and heavy rainfall, with 24 hour accumulations of greater than three inches. This combination often results in moisture entering buildings, creating conditions for the growth of mold. Examining the health effects of specific contaminants in buildings is very complex, but research has shown that some respiratory symptoms and illnesses can be associated with damp buildings.<sup>47</sup> According to County Health Rankings, 18 percent of households in Lincoln County have severe housing problems.

## **Tobacco-free Spaces**

Tobacco use is still the leading preventable cause of death and disability in Lincoln County. Statistics on tobacco related diseases and deaths are discussed in Chapter 6: Morbidity and Mortality.

Tobacco use is directly linked to coronary heart disease, stroke, cancer and chronic lung disease. Tobacco use is also linked to additional health and environmental concerns such as exposure to second hand smoke; fire related death and injury; increased risks of wildfires; and littering of toxic cigarette filters. In order to reduce these health and environmental impacts, Lincoln County and the state have taken steps to reduce exposure to tobacco and cigarette smoke in public places.

Currently, Oregon law prohibits smoking and other tobacco products in most workplaces, schools, bars, and other indoor public spaces. Lincoln County allows smoking on county property at some county parks and waysides, but has restricted it on all other county-owned property. Furthermore, many governmental bodies are expanding smoke- or tobacco-free policies to explicitly include e-cigarettes.<sup>48,49,50</sup>

Within the region, number of non-governmental entities also restrict or ban tobacco on their properties. Oregon Coast Community College and Samaritan Health Services are among the entities that ban tobacco products on their property.<sup>51,52,53</sup>

## **Transportation**

Transportation links people and places, making it possible to get to work, to school, to recreational opportunities, and to the grocery store. Transportation includes more than roads, walkways, or bridges. It also encompasses public transit systems, policies that dictate the location and construction of roads, and guidelines for accommodating different kinds of users. Guidelines are important for providing avenues for physical activity, and for reducing the potential of driver, cyclist, and pedestrian injury.

## Access to Public Transportation

Access to public transportation is an important public good. Not only does taking public transportation provide additional opportunities for exercise, but the presence of public transportation also makes it easier for individuals and families without private transportation to access goods and services vital to maintaining health. These include grocery stores, health and dental care, and recreation facilities. In Oregon, counties with large metropolitan areas relative to county population size tend to have more public transportation options. Lincoln County's large land area and the greater distances between its small cities and towns pose challenges for county and city public transit. Twenty-nine percent of Lincoln County residents live within one quarter of a mile of a bus stop (ranked 8<sup>th</sup> in the state).<sup>54</sup> Although distance to a public transportation route is one measure of the strength of a public transportation system, additional factors impact the strength of public transport, including frequency and hours of operation, direct routes, and connections to other routes.

People of color, people experiencing poverty, people with disabilities, and people who experience language barriers are more likely to depend on public transit. However, they often live in areas with poor transit service, fewer destinations, and poor connectivity. These unfair burdens increase transportation costs and stress, and limit access to economic and educational opportunities, housing, healthy foods, and physical activity. Vulnerable populations often have unsafe transportation conditions, including limited safe crossings, areas with high-speed traffic, and poor sidewalk and bicycle infrastructure.

## Access to Healthy Foods

Transportation options and limited public transportation for Lincoln County residents contributes to challenges with regard to nutritious food access. For households without private vehicles, the ability to shop for food at grocery stores is highly dependent on proximity. Twenty-three percent of households in Lincoln County are within one half mile of a grocery store. The average distance between a household and the nearest grocery store is 2.6 miles.<sup>55</sup> However, since grocery stores tend to be located in larger towns, the county average may overestimate the urban average and underestimate the rural average.

Access to nutritious foods can be particularly difficult for residents with unreliable transportation or tight budgets. A rural community is considered to have low access to food when it is ten or more miles from a supermarket or large grocery store. Rural residents must often travel long distances for food. Rural grocery stores throughout the county report barriers that may limit rural low-income families' access to healthy food. These include: administrative barriers to becoming an authorized vender for SNAP and WIC programs, economic barriers to offering fresh fruits and vegetables, meat, dairy and other refrigerated foods.<sup>56</sup>

As a comparison, many more Lincoln County residents live within one half mile of a tobacco vendor compared to within one half mile of a grocery store (Table 3.1). Approximately six percent of Lincoln County residents are low income and do not live close to a grocery store.<sup>1,57</sup>

Table 3.1: Proximity to grocery stores compared to tobacco vendors in Lincoln County, 2012

Vendor	Average (mean) walking distance in miles	Percent of population living within ½ mile
Grocery stores	2.6	23 %
WIC-authorized stores	2.7	19 %
Tobacco vendors	1.7	37 %

Source: Oregon Environmental Public Health Tracking, 2012

In addition to access to nutritious food, proximity to fast food can affect the health of the community. Although complex in nature, the food environment can impact what people eat, and providing healthy options is vital for the health of the community. Although not causal, studies have shown an increase in the prevalence of obesity and diabetes with increased access to fast food outlets in a community. Twenty-one percent of restaurants in Lincoln County are fast food vendors.<sup>58</sup>

## Conclusion

The previous Community Health Needs Assessment did not include a section on the environment, and changes to the environment generally take place over long time frames. Therefore there are no comparisons that can be made with the previous CHNA.

---

<sup>1</sup> “Close” is defined as within 1 mile for urban areas and within 10 miles for rural areas



# Chapter 4

## Social Determinants of Health

Opportunities for health among residents of Central Lincoln County begins within their communities including their homes, neighborhoods, places of worship, workplaces, and schools. A growing body of scientific research shows that all people benefit when communities invest in health.

The World Health Organization defines social determinants of health as “the conditions in which people are born, grow, work, live and age, and the wider set of forces and systems shaping the conditions of daily life”.<sup>59</sup> These non-medical factors contribute to a large percent of preventable poor health outcomes. Social determinants include influences such as: “early years’ experiences, education, economic status, employment and decent work, housing and environment, and effective systems of preventing and treating ill health.”<sup>60</sup> These aspects of health are often referred to as “upstream factors” since their effect occurs well before illness manifests and curative intervention becomes necessary. In this chapter Central Lincoln County data will be presented for education, employment, income, poverty, economic challenges, food security, home ownership, and homelessness. Where Central Lincoln County data is not available, Lincoln County data is reported.


### Income, Poverty, and Economic Challenges

Income is the strongest predictor of health among all social determinants of health. Not only are there many studies showing a strong association between income and health<sup>61</sup>, but income also affects all other social determinants of health, including education, food security, and housing. The National Longitudinal Mortality Survey found that people in the top five percent of incomes had life expectancies 25 percent longer than people in the bottom five percent of incomes.<sup>62</sup> While income is not a “one size fits all” measure of health, understanding the income characteristics of Central Lincoln County provides a solid foundation for measuring social determinants of health in Central Lincoln County.

#### Median Incomes

The median income of a population is a measure of the average income in that population; 50% of the population earns more than the median income, and 50% of the population earns less. The median (inflation-adjusted) household income for Central Lincoln County ranges between \$36,000 and \$53,000 depending on the Census Tract; the median household income for all of Lincoln County is \$42,395.

Figure 4.1: Per capita inflation-adjusted income by subpopulation group in Lincoln County, 2011-2013.



Source: U.S. Census Bureau American Community Survey, 2010-2014

## Income inequality

Income inequality (the distribution of wealth between richer and poor segments of the population) is associated with many health outcomes. Regions with higher inequality are more likely to experience increased infant mortality, lower life expectancy, higher rates of depression, and lower health status overall. Income inequality is commonly measured by calculating the ratio of the 80<sup>th</sup> income percentile to the 20<sup>th</sup> income percentile of the population.<sup>263</sup> In Oregon, the 80<sup>th</sup> income percentile is 4.6 times the 20<sup>th</sup> income percentile. Central Lincoln County has a ratio of 4.5, slightly lower than the state ratio.

## Poverty

Poverty is inextricably linked to poor health outcomes. Poverty is related to both limited income and lack of economic stability, limited choices in education, employment, and living conditions, and reduced access to safe places to live, work, and play. It can also frequently hinder choices and access to healthy food.

The United States Census Bureau determines the Federal Poverty Level (FPL) each year. The FPL was originally an estimate of the amount of money required to meet the cost of living for individuals or families. Currently, the FPL is a statistical threshold of poverty.<sup>64</sup> It is not generally recognized as an accurate measure of true poverty, but it is used for determining

<sup>2</sup> The 80<sup>th</sup> income percentile is the income of the individual who earns more than 80 percent of the population. The 20<sup>th</sup> income percentile is the income of the individual who earns more than 20 percent of the population. Those who earn more than the 80<sup>th</sup> income percentile are the richest 20% of the population; those who earn less than the 20<sup>th</sup> percentile are the poorest 20% of the population.

eligibility for assistance programs. Below, in Table 4.1, the FPL for individuals and families is presented, as well as specific FPL ratios that are used for eligibility and comparison purposes.


Table 4.1: Annual Income and Federal Poverty Levels and related ratios for 2013.

Family size	Percent of Federal Poverty Level					
	50%	<b>100%</b>	138%	185%	200%	400%
<b>Individual</b>	\$5,940	<b>\$11,880</b>	\$16,394	\$21,978	\$23,760	\$47,520
<b>Three person family</b>	\$9,277	<b>\$18,554</b>	\$25,605	\$34,325	\$37,108	\$74,216
<b>Four person family</b>	\$11,922	<b>\$23,844</b>	\$32,905	\$44,111	\$47,688	\$95,376

Source: U.S. Census Bureau, Historical Poverty Threshold Table

Approximately 16 percent of Central Lincoln County’s population lives below the federal poverty line, compared to 17 percent of Oregon’s total population. One worrisome statistic is that children less than 18 years of age are among the age groups with the highest percentage living below the federal poverty level in Lincoln County.<sup>65</sup> 26 percent of children less than 18 years of age are below the poverty level.

Figure 4.2: Percent of population living below the federal poverty line by age group in Lincoln County, 2010-2014




Source: U.S. Census Bureau, American Community Survey, 2010-2014

Earning less than a high school education increases the risk of experiencing poverty.<sup>66</sup> In Lincoln County, 25 percent of adults over the age of 25 who did not complete high school are below the federal poverty line, compared with 12 percent of those who completed high school.

Variation in poverty rates also exists between racial/ethnic groups in Lincoln County. As shown in Figure 4.3, all racial and ethnic groups in Lincoln County have a higher poverty rate than the White, non-Hispanic/Latino population except for Other. Individuals in Lincoln County who identify as Hawaiian or Pacific Islander and Black or African American are among the

race/ethnic groups with the highest poverty rates with 87 percent and 28 percent, respectively.<sup>67</sup> It is important to note, however, that the populations for non-White and non-Hispanic/Latino groups are very small.

Figure 4.3: Percent of population living below the federal poverty line by race and ethnicity, 2010-2014



Source: U.S. Census Bureau, American Community Population, 2010-2014

## Low Income and Cost of Living

Many Lincoln County residents earn incomes higher than the federal poverty level but still struggle economically to meet their everyday needs. Nearly 31 percent of Lincoln County earns less than 185 percent of the federal poverty level (\$21,775 annually for an individual or \$44,863 annually for a family of four in 2015).<sup>68,69</sup> This is the threshold that many assistance programs, such as the Supplemental Nutrition Assistance Program (SNAP), use for income eligibility.

Research suggests that the cost of living in Lincoln County is well above the federal poverty level. Table 4.2 below shows the cost of living for three family types in each county, and the corresponding poverty level. These figures take into account costs such as housing, child care, food, transportation health care, and taxes.

Table 4.2: Cost of living as a percent of the federal poverty level, 2014

County	One adult, one preschooler		One adult, one preschooler, one school-age		Two adults, one preschooler, one school-age	
	Annual cost of living	Annual cost of living as percentage of FPL	Annual cost of living	Annual cost of living as percentage of FPL	Annual cost of living	Annual cost of living as percentage of FPL
<b>Lincoln</b>	\$32,390	206%	\$36,624	185%	\$45,918	193%

## Employment

Stable and secure employment influences health, not only by being a source of income, but also by providing access to health insurance. Compared to unemployed workers, individuals who are employed fulltime have higher incomes and standards of living, less stress, and may be less likely to turn to unhealthy coping behaviors such as alcohol consumption or smoking.<sup>70</sup> The unemployment rate has been decreasing steadily in recent years.<sup>71</sup> As of May 2016, the seasonally adjusted unemployment rate in Lincoln County was 5 percent.<sup>72</sup> Generally, an unemployment rate of 5% is considered “full employment” as there is always a certain amount of turnover in the labor force.

## Economic Opportunities

Lincoln County’s top business sector comprises the arts, entertainment, recreation, accommodation and food services (21 percent). The second biggest business sector in Lincoln County is the education, healthcare, and social assistance sector, which employ 19 percent of the working population. Other important business sectors across in Lincoln County are retail trade; manufacturing; and professional, scientific, management, administrative, and waste services.<sup>73</sup>

## Education

Health and education are closely connected. Educational access and attainment are very important predictors of health status. Individuals with higher levels of education are less likely to die prematurely or report acute diseases. They also report positive health behaviors, like maintaining healthy weight, and fewer risky behaviors, like smoking.<sup>74</sup> Furthermore, education levels are the strongest predictor of income and wealth, which strongly influence lifelong health.<sup>75</sup>

## Early Learning

Early childhood development supports nurturing relationships and learning opportunities that foster children’s readiness for school. The early years are crucial for influencing health and social well-being across a child’s lifetime. Research evidence accumulated over the past 40 years supports the conclusion that children who participate in high-quality early childhood development (ECD) programs benefit from a broad range of immediate and long-term health benefits.<sup>76</sup>

The Head Start Program is one such federal program that promotes the school readiness of children from low-income families by enhancing their cognitive, social, and emotional development. Head Start programs provide a learning environment that supports children’s

growth from birth to age five in several areas, such as language, literacy, and social and emotional development. Head Start programs also emphasize the role of parents as their child’s first and most influential teacher, and support the development of healthy familial relationships and well-being.<sup>77</sup> In Oregon, Head Start programs include the Oregon Head Start Prekindergarten (OHS PreK) program, which serves children age three to five from low-income families. Some Head Start programs also include Early Head Start (EHS), which is a comprehensive program for children below the age of three and pregnant women from low-income families. Oregon children whose families are below the federal poverty level (\$24,250 for a family of 4) are eligible for these benefit programs.<sup>78-</sup>

The OHS PreK and EHS programs that serve children and families in Lincoln County are shown in Table 4.3 below:

Table 4.3: Oregon Head Start PreK and Early Head Start programs and enrollment, 2013-2014

OHS PreK and EHS program	County	OHS PreK enrollment	EHS enrollment	Total enrollment
<b>Community Services Consortium</b>	Lincoln	154	--	154
<b>Siletz Tribal Head Start</b>	Lincoln	20	30	50
<b>Total</b>		174	30	174

Source: Oregon Department of Education, Early Learning Division, Oregon Head Start Prekindergarten Programs 2013-2014 Directory

Despite strong research showing the positive impact of high-quality early education, many families in Lincoln County who are in need of child care may not be served. While there are not data for informal child care options, in 2012, for every 100 children there were 17 available child care slots in Lincoln County. In Oregon, there were also 17 available child care slots per 100 children. The goal for the state is 25 slots per 100 children; Lincoln County has not yet met that goal. In addition to availability, price may be a barrier for many families. The average annual cost of toddler care in childcare centers in Lincoln County is \$6,240, is lower than the average cost in Oregon.<sup>79,80,81</sup>


## High School Education

High school graduation is a strong predictor of future employment and earnings. Conversely, dropping out of school is associated with lower income, multiple social and health problems,<sup>82</sup> and health risks.<sup>83</sup> For example, 32 percent of Oregonians who do not have a high school degree smoke, compared with 24 percent of high school graduates, 18 percent with some post-secondary education, and seven percent of college graduates (age-adjusted).<sup>84</sup>

In the 2013-2014 school year, Lincoln County experienced a high school dropout rate of 47 students per 1,000 9<sup>th</sup>–12<sup>th</sup> graders. In general, Lincoln County’s dropout rates were comparable with the state dropout rate between 2008 and 2014. Central Lincoln County had a high school graduation rate of 71%.

In 2011, Oregon set a goal of 40-40-20, meaning that by 2025, 40 percent of Oregonians would have a bachelor’s degree or higher, an additional 40 percent would have an associate’s degree, and the remaining 20 percent would have graduated high school. This translates to 100% of Oregonians having a high school degree or higher, and 80% having an associate’s degree or higher. In 2013, 89% of Central Lincoln County residents had completed high school or GED equivalent, 33% had an associate’s degree or some college, and 24% had a bachelor’s degree or higher (Figure 4.4).<sup>85</sup>

Figure 4.4: Rates of high school completion, associate’s degree or higher, and bachelor’s degree or higher, In Lincoln County, 2015



Source: County Health Rankings 2015


## Food Security

Food security is defined as having enough to eat, and being able to purchase or obtain healthy food in socially acceptable ways.<sup>86</sup> Adequate nutrition is particularly important for children, as it affects their cognitive and behavioral development. Children from food insecure, low-income households are more likely to experience irritability, fatigue, and difficulty concentrating on tasks, especially in school, compared to other children.<sup>87</sup>

Feeding America, a national nonprofit that monitors food security, estimates that 27 percent of children in Lincoln County are living in food insecure households as shown in Figure 4.5 below.



Figure 4.5: Percent of children living in food insecure households in Lincoln County, 2013



Source: Feeding America

Based on Oregon Department of Education data, 68 percent of Lincoln County K-12 students were eligible for free/reduced lunch during the 2013-2014 school year. Students whose family incomes are below 130 percent of the federal poverty level (\$31,525 annually for a family of four) are eligible for free lunches, and students whose family incomes lie between 130 and 185 percent of the federal poverty level (between \$31,525 and \$44,863 annually for a family of four) are eligible for reduced-price lunches.<sup>88,89</sup>

An analysis of factors<sup>3</sup> determining food insecurity suggests that in 2013, 15 percent of Lincoln County population, approximately 7,000 individuals, were residing in households that were food insecure. Among those who were food insecure, 21 percent earned incomes above 185 percent of the federal poverty level, making them ineligible to participate in government assistance programs (Table 4.4). The childhood food insecurity rate was much a higher, at 27 percent of the children in Lincoln County. Of the children living in food insecure households in Lincoln County, it is estimated that 27 percent of these children are likely ineligible for federal nutrition programs because they live in households with incomes above 185 percent of the federal poverty level.<sup>90</sup>

Table 4.4: Food insecurity in Lincoln County and Oregon, 2013

	Number of food insecure individuals	Percent of population that is food insecure	Percent of food insecure people who are ineligible for benefits *
<b>Lincoln County</b>	7,070	15%	21%
<b>Oregon</b>		27%	27%

\* Percent ineligible figure is produced by modeling and is an estimate

Source: Feeding America

<sup>3</sup> Factors include indicators of food insecurity such as poverty, unemployment, median income; food budget shortfalls; a cost of food index; and national average meal costs.

## Supplemental Nutrition Assistance Program Participation

The Federal Supplemental Nutrition Assistance Program (SNAP) is the largest domestic food and nutrition assistance program for low-income Americans. U.S. households must meet certain eligibility criteria, such as income, to receive benefits. In 2013, it is estimated that 22 percent of all households (1,201 households) in Central Lincoln County received SNAP benefits.

## Women, Infants and Children (WIC)

WIC is a public health nutrition program that is vital to the health of women, infants, and children across Oregon. The WIC program provides health and nutrition services to pregnant and breastfeeding women and children ages 0 to 5 that have a household income less than 185 percent of poverty guidelines.<sup>91</sup> Overall in 2013, a total of 2,260 families were served by WIC in Lincoln County; 71 percent of these were infants and children under five, and 29 percent were pregnant, breastfeeding, and post-partum women. Approximately 62% of pregnant women in Lincoln County were served by WIC. Furthermore, 71 percent of families served by WIC in Lincoln County were working families.<sup>92,93,94</sup>

## Emergency Food Support

The Food Share of Lincoln County provides food boxes, a food pantry, and emergency meals. Between July 2014 and July 2015, the Lincoln County Food Share distributed enough food to feed approximately 400 people three meals a day for a year.<sup>95</sup>

## Housing and Home Ownership

Housing is an important part of the built environment and another key factor contributing to good health. Older housing in particular can present multiple threats to health, including the presence of mold, asbestos, lead-based paint, and lead solder in plumbing and in the soil.

Poor quality and inadequate housing contribute to health problems such as infectious and chronic diseases, injuries, and poor childhood development. Indoor allergens and damp housing conditions play an important role in respiratory conditions including asthma, which currently affects over 20 million Americans and is the most common chronic disease among children. Approximately 40 percent of diagnosed asthma among children is believed to be attributable to residential exposures.

## Housing Affordability

Affordable, quality housing provides shelter that is safe and healthy for all people. Housing that costs more than 30 percent of household income is considered to be “unaffordable.”<sup>96</sup> Thirty-six percent of South Lincoln households are cost burdened.

## Homelessness

The Oregon’s Ending Homelessness Advisory Council defines homelessness as being without a decent, safe, stable, and permanent place to live that is fit for human habitation.<sup>97</sup>

Understanding homeless populations is a daunting challenge for public health. Homeless people are just as much a part of society as housed individuals, but they face additional obstacles in accessing social services or health care. Even counting the number of homeless individuals is a difficult task, because a homeless individual may move around a lot during the year or be unwilling to interact with social services. Each January, Oregon Housing and Community Services require communities to conduct a point-in-time count of homeless populations. This snapshot of the homeless population is limited in scope and depth. Canvassers visit shelters, transitional housing, and known homeless encampments. Individuals staying with other people out of economic necessity are not counted, nor are homeless people who are in areas not covered by the canvassing. Furthermore, the one-night count misses any individual who is homeless at other points during the year. Notwithstanding these limitations, the point-in-time estimates have the benefit of being a consistent approach across years and geographies, and therefore may give some insight into the homeless community in Lincoln County.

In 2011, the county point-in-time surveys counted 41 homeless individuals in Lincoln County (Table 4.5). All of these individuals were in shelters or transitional housing. There were no street counts conducted in Lincoln County in 2011. Forty percent of the homeless population in Lincoln County was male. The average length of time spent homeless was 20 months for men and 7 months for women.

The most recent data on homeless populations is from 2015 (Table 4.7). In 2015, there were 47 homeless individuals identified in the January point-in-time survey. However, in 2015 the unsheltered counts comprised a larger portion of the records, which may indicate a larger canvassing effort rather than solely an increase in the homeless population.

In both 2011 and 2015, approximately one quarter of the recorded individuals were members of families, both adults and children (Table 4.5).

Table 4.5: One-night count homeless population figures in Lincoln County

	2011	2015
<b>Total homeless count</b>	41	47
<b>Sheltered count</b>	41	25
<b>Unsheltered count</b>	0	22
<b>Male</b>	16*	23
<b>Female</b>	24*	23
<b>Individuals</b>	23	29
<b>Family members</b>	18	18
<b>Average months spent homeless (male / female)</b>	20/7	No data

\* Counts do not sum to total

Source: OHCS and Community Services Consortium

Another source for recording the number of homeless individuals is the set of statistics gathered by federally qualified health centers (FQHCs). Among the data that FQHCs are required to collect is housing status, which they report each year to the federal government. According to the Bureau of Primary Health Care, a patient’s status should be recorded as homeless if the patient was residing in a shelter, transitional housing, on the street, if the patient was doubled up or temporarily living with others, had been homeless within the last 12 months, or resided in a housing program targeted to homeless populations. Compared with the one-night counts, FQHCs may identify homeless individuals who were not staying in shelters or in canvassed encampments or who were homeless at other times throughout the year. However, only those individuals who were able to seek out medical care at an FQHC and chose to do so were identified. Nevertheless, the records provided by the FQHCs indicate a much broader level of homelessness than the one-night counts. In 2014, the Lincoln FQHC served approximately 564 homeless patients who were homeless during at least one visit to the FQHC. This number is twelve times as large as the 2015 one-night count, and represents a 50 percent increase from the number of homeless FQHC patients in 2012.<sup>98</sup>

Student homelessness is a recurring problem in Oregon as well. Across the state, an increasing number of Oregon’s K-12 public school students are homeless at some point during the school year. Homelessness among students has more than doubled since the 2003-2004 academic school year. 10 percent of Central Lincoln County K-12 students experienced homelessness in the 2014-2015 academic year (519 students).

## Conclusion

The major change in the socioeconomic status of Lincoln County is the unemployment rate, decreased from 9.5 percent in 2012 to 5 percent in 2016. Other comparable indicators have stayed largely the same.

# Chapter 5

## Access to Medical Care

It is important to examine medical care access and capacity in the larger context of overall factors that contribute to health. “Health care is necessary but not sufficient for improved health; in fact, health care accounts for only about 10–20 percent of health outcomes, according to some experts.”<sup>99</sup> Social determinants of health, the upstream factors listed in the previous chapter, are responsible for a much larger percentage of health outcomes than medical care alone. People need a healthy and accessible environment to achieve good health. This includes the broader community context, as well as the characteristics of the local health care system itself.

Many of the forces that shape the opportunity for better health in Central Lincoln County – education, employment, and transportation, for instance – can also affect access to medical care. Hospital data is reported for Samaritan Pacific Communities Hospital; other data is reported at the county level.

### Demographic Differences in Access to Medical Care

Some populations face increased barriers to accessing care and receive poorer quality care when they get it. In its 2011 reports on health care quality and disparities, the Agency for Healthcare Research and Quality (AHRQ) finds that, at a national level, low income individuals and people of color experience more barriers to care and receive poorer quality care. Moreover, other research shows that individuals with limited English proficiency are less likely than those who are English proficient to seek care even when insured. Research also finds differing patient experiences and levels of satisfaction by race, gender, education levels, and language.<sup>100</sup>


### Health Insurance Coverage

Lack of adequate health insurance coverage is often a major barrier to medical care. People who are uninsured or underinsured receive less medical care than their insured counterparts.<sup>101</sup> Inadequate coverage creates a financial barrier between a patient and needed medical care. People without health insurance are less likely to know about or seek out preventive services, and are more likely to have new and worsening health problems, and shorter lifespans.<sup>102</sup> In general, even when uninsured/underinsured persons receive medical care, care is often postponed (due, in part, to concerns about cost). As part of the ACA, Oregon accepted federal funding to expand Oregon Health Plan (OHP) membership, setting targets for enrollment and expanding the variety of services (e.g. dental services). Statewide, membership in OHP increased 75 percent over four years, from 469,000 members in January 2010 to 821,000

members in January 2014. Lincoln County enrollment increased from 6,000 members to 12,000 members over the same time period.<sup>103,104</sup> In addition to OHP expansion, eighty percent of the consumers registered to the new health care exchange received tax credits and/or cost-sharing subsidies as of April 2014.<sup>105</sup>

Insurance coverage rates in Lincoln County have risen recently, largely due to the ACA and other healthcare transformation policies. Lincoln County insurance coverage rate in 2012 was 82 percent, rising to 96 percent in 2014.<sup>106</sup> (Figure 5.1).<sup>107</sup>

Figure 5.1. Insurance coverage rates over time in Lincoln County



Source: U.S. Census Bureau, American Community Survey, 2009-2013

\* 2014 coverage rates are from Oregon Health Authority, 2015

Because of the rapidly shifting health care and health insurance landscape, local data points that accurately capture these changes are still forthcoming. With that in mind, data from before the ACA expansion showed major disparities among the population based on age, race, and income. Examining these disparities across Lincoln County can help provide a baseline for future comparisons with disparities which exist after ACA expansion once the data is available.

## Uninsured rates

Uninsured rates differed greatly between age groups before ACA. The uninsured rate among children across Lincoln County was lower than the rate for working-age adults (Table 5.1).<sup>108</sup> Across Lincoln County, less than one percent of individuals 65 and older lack health insurance due to Medicare. The age group with the highest uninsured rates in Lincoln County was 18 to 24 year olds, at 28 percent.

Table 5.1: Uninsured rates in Lincoln County, 2011-2013

Lincoln County	
<b>Under 18 years old</b>	11.4 %
<b>18 to 64 years old</b>	28 %
<b>65 years old and older</b>	0.33 %

Source: U.S. Census Bureau, ACS 2011-2013, Table S2702

Insurance coverage rates were also pronounced across racial/ethnicity categories, employment status, and citizenship status. From 2009 to 2013 in Lincoln County, over 34 percent of Latino individuals and over 30 percent of American Indian and Alaska Native were uninsured, compared to 17 percent of the White population. Black or African American and two or more races had an uninsured rate of 13 percent. Additionally, 52 percent of the unemployed are uninsured, compared to 24 percent of those currently employed. The foreign born and non-citizens have very high uninsured rates, at 50 percent and 63 percent, respectively.<sup>109</sup> Insurance coverage data is not available for undocumented immigrants. However, undocumented immigrants, including undocumented children, are excluded from both Medicaid and the health insurance exchange.<sup>110</sup>

Among the employed, those working less than full time year-round were uninsured at a higher rate (39 percent) compared to those working full time year-round (20 percent). Residents earning less than 200 percent of the federal poverty level are more likely to be without insurance coverage than those with higher incomes, 30 percent versus 12 percent.<sup>111</sup> The implementation of the Affordable Care Act has had a major impact on insurance coverage rates in Lincoln County, as Figure 5.1 demonstrates. However, even given the growth in insurance coverage rates over the past 5 years, insurance gaps and inequalities remain, especially for people of color, individuals living in rural areas, and low income workers.<sup>112</sup> As data for recent years become available, it will be important to measure these disparities.

## Health insurance among children

Examining insurance coverage rates among children up to age 18 shows a gradual increase in Lincoln County from 2006 to 2012. As of 2012, Lincoln County had an insurance coverage rate of 92 percent for children under the age of 18. While data are not yet available to demonstrate the effects of the ACA on insurance coverage rates among children, the upward trend preceding the ACA expansion provides an important baseline.

## Cost of Medical Care

Insurance coverage is only part of the cost of medical care. Additional costs are referred to as cost-sharing and include costs such as copayments, coinsurance and deductibles. Health reform legislation has reduced financial burdens for many people with lower income or significant health care needs. Nevertheless, one in three Americans say they have put off getting medical treatment that they or their family members need because of cost.<sup>113</sup>

According to the County Health Rankings, during the 2006-2012 period, 19 percent of adults in Lincoln County reported they did not see a doctor in the past 12 months because of cost.<sup>114</sup>

## Access Capacity

Primary care, mental health, and oral health are foundational to a comprehensive offering of medical care for a population.

Other primary care providers are especially vital in rural areas that may not have the population density to support a full time physician.

Lincoln County had 1,775 residents per primary care physician; 784 residents per primary care provider; 400 residents per behavioral health provider, and 1,799 residents per oral health provider.

Having a usual primary care provider (PCP) is associated with improved health outcomes, increased health equity, and lower healthcare costs. Effective PCPs work to maintain sustainable relationships with patients, connect them with additional health resources in the community, and coordinate their care. Patients with ongoing access to PCPs and other healthcare services have better relationships with their providers and are more likely to receive appropriate care than patients without a regular healthcare provider.<sup>115</sup>

## Safety Net Services & Community Benefits

The health care “safety net” refers to the component of the health care system serving low-income and uninsured people. Safety net services are complemented by community funding, programs and activities.<sup>116</sup>

Federally Qualified Health Centers<sup>4</sup> (FQHCs) and Free Clinics or “charity” clinics are the most common types of safety net clinics. FQHCs in Lincoln County provide primary care, mental/behavioral health, and oral health services. Central Lincoln County has an FQHC located in Newport. Additionally, there are school-based health centers acting as FQHCs in Newport, Toledo, and Waldport.<sup>117</sup> In Lincoln County, FQHCs served just under 5,000 patients in 2014.<sup>118</sup> Auxiliary safety net providers also serve the region’s vulnerable populations. Persons served by these programs include homeless, women and children through maternal-child and WIC services, and HIV-positive people.

---

<sup>4</sup> FQHCs have a legal mandate or expressly adopted mission to serve all patients, regardless of ability to pay or legal status.



## Health Care Professional Shortage Areas

Knowing the number of providers and types of services are very important for gauging the capacity and presence of a health care system. However, an understanding of the geographical distribution of these services helps paint a more accurate picture.

While Lincoln County enjoys a good ratio of health care providers to overall population, geographic distribution of providers can make it difficult for those with limited transportation to access services. Because rural areas of Lincoln County have either no or very few medical care providers, portions of Lincoln County are designated as geographic Health Care Professional Shortage areas (HPSA). Designation as an HPSA means that there is an increased risk of poor access to health professionals.<sup>119</sup> Lincoln County qualifies in part as an HPSA for primary care, dental health, and mental health.

In addition to the geographic designation, Lincoln County also has population-based HPSAs for migrant seasonal farmworkers and low income individuals. Migrant seasonal farmworkers and their families are a particularly vulnerable subgroup of the Latino/Hispanic population. Farmworkers have different and more complex health problems than those of the general population. Many of the Latino/Hispanic migrant seasonal farmworkers are documented but have undocumented family members with them. Many are employed in agriculture sectors that provide few or no employment benefits. While most are low income, many immigrants and migrant seasonal farm workers do not qualify for Medicaid due to their residency status or they are unable to access Medicaid due to language, transportation and cultural barriers.<sup>120</sup>

## Emergency Responders

Emergency Management Services (EMS) serve an important role in the community. According to the Oregon Office of Rural Health, the mean travel time to the nearest hospital for rural service areas is 24 minutes. Estimated travel time is calculated from the largest town/city in each of the rural service areas to the nearest town/city with a hospital. This is the protocol unless the city already has a hospital, in which case driving time is defaulted to 10 minutes.<sup>121</sup> The service area covered by Samaritan Pacific Communities Hospital is all within 24 minutes by car.<sup>122</sup>

## Medical Services

The following eight indicators provide a snap-shot of the breadth and type of services provided by Samaritan Pacific Communities Hospital (SPCH) over the past three years:

- The number of inpatient visits
- The number of surgeries performed
- The number of infants delivered
- The number of imaging procedures performed


- The number of ER visits
- The number of clinic visits
- The number of hospice visits
- The number of home health visits

## Inpatient Visits

Inpatient care begins when a doctor makes a formal order to admit a person as an inpatient. The length of inpatient care depends on the severity of the health issue and when the doctor deems it safe for the patient to leave.

From 2013-2015, there were on average 1,300 inpatient visits each year at SPCH. The annual rate decreased 9 percent from the previous triennium, when on average 1,440 visits were made annually. The following figure (Figure 5.2) illustrates the six year trend.

Figure 5.2: Annual inpatient visits, 2010 - 2015




Source: Samaritan Health Services, 2016

## Surgical Procedures

Samaritan Pacific Communities Hospital offers surgical services in a number of specialties, including inpatient, outpatient, and minimally invasive surgeries using the best technology available.<sup>123</sup>

From 2013-2015, there were on average 2,290 surgical cases each year at SPCH. The number of surgical procedures increased five percent over the last triennium.

Figure 5.3: Annual surgical cases, 2010 - 2015




Source: Samaritan Health Services, 2016

## Infant Deliveries

The Mark G. Collson Birthing Center at SPCH offers an array of programs and services to women from all backgrounds.<sup>124</sup>

From 2013-2015, there were on average 179 infant deliveries each year at SPCH, down from 205 in the previous triennium. Deliveries have been declining steadily over the last six years, from 222 in 2010 to 171 in 2015. The following figure (Figure 5.4) illustrates the six year trend.

Figure 5.4: Annual infant deliveries, 2010 - 2015



Source: Samaritan Health Services, 2016


## Imaging Procedures

SPCH offers a variety of imaging procedures:

- Bone Density Test (DEXA)
- Digital Mammography
- Echocardiography
- MRI
- Ultrasound (2-D)
- X-ray
- X-ray (Digital)<sup>125</sup>

SPCH has been increasing the number of imaging procedures it conducts. Between 2013 and 2015, SPCH conducted 30,450 imaging procedures annually, a 17 percent increase over the previous triennium. The following figure (Figure 5.5) illustrates the six year trend.

Figure 5.5: Annual imaging procedures, 2010 - 2015




Source: Samaritan Health Services, 2016

## Emergency Department Visits

People without health insurance and access to other providers often end up seeking emergency room care. Uninsured adults are more likely than those with private or public health insurance to utilize emergency room care because they lack preventative care and places to receive affordable, accessible care. Lack of access to other providers and adequate health insurance is reflected in rising visits to the emergency room.<sup>126</sup>

From 2013-2015, there were on average 15,700 emergency department visits each year at SPCH, an increase of 22 percent over the last triennium. Emergency department visits have increased by nearly fifty percent over the past six years. The following figure (Figure 5.6) illustrates the six year trend.

Figure 5.6: Annual emergency department visits, 2010 - 2015



Source: Samaritan Health Services, 2016

## Medical Clinics

There are twelve clinics serving the SPCH area. The following table lists each clinic along with its area of practice and location.


Table 5.2 Medical clinics in the SPCH service area

Clinic Name	Area of Practice	Location
<b>Samaritan Depoe Bay Clinic</b>	Family Medicine, Walk-in Clinic, Specialty Services	Depoe Bay
<b>Samaritan Hematology &amp; Oncology Consultants</b>	Specialty Services	Newport
<b>Samaritan Kidney Specialists</b>	Specialty Services	Newport
<b>Samaritan Occupational Medicine</b>	Specialty Services	Newport
<b>Samaritan Orthopedic Surgeons</b>	Specialty Services	Newport
<b>Samaritan Pacific ENT &amp; Allergy</b>	Specialty Services	Newport
<b>Samaritan Pacific Internal Medicine</b>	Internal Medicine	Newport
<b>Samaritan Pacific Surgical Associates</b>	Specialty Services	Newport
<b>Samaritan Pacific Walk-in Clinic</b>	Walk-in Clinic	Newport
<b>Samaritan Pacific Women’s Health Group</b>	Obstetrics/Gynecology	Newport
<b>Samaritan Toledo Clinic</b>	Family Medicine	Toledo
<b>Samaritan Waldport Clinic</b>	Family Medicine, Pediatrics	Waldport

Source: Samaritan Health Services, 2016

From 2013-2015, there were on average 55,350 clinic visits each year within the SPCH service area, slightly more than in the previous triennium. However, in the last three years, clinic visits have increased by over 33 percent, from 48,000 in 2013 to 65,000 in 2015. The following figure (Figure 5.7) illustrates the six year trend.

Figure 5.7: Annual clinic visits, 2010 - 2015




Source: Samaritan Health Services, 2016

## Hospice Care

Samaritan Pacific Hospice offers long-term care resources for patients who are nearing end of life. Hospice services support patients emotionally, spiritually, and physically during this transition. Families of patients are also provided emotional support during this period of grief.<sup>127</sup>

Hospice utilization is measured by the total number of days of hospice care provided. From 2013-2015, the annual hospice days varied greatly. In 2013-2015 there were on average 3,800 hospice days each year 12 percent than in the previous triennium. However, the number of hospice days each year varies by up to 1,000 days from year to year. The following figure (Figure 5.8) illustrates the six year trend.

Figure 5.8: Annual hospice days, 2010 - 2015



Source: Samaritan Health Services, 2016


## Home Health Care

Patients recovering from a stroke, accident, or coping with a long-term illness may require medical assistance within the home. These services often include physical therapy, home medical equipment assistance, coping skills education, diabetes management, and other resources designed to maintain or improve a patient's health at home.

Each year a variety of health professionals visit patients in their homes. At SPCH, this includes, registered nurses (RN), physical therapists (PT), occupational therapists (OT), speech therapists, medical social worker visits (MSW) and home health aides.

From 2013-2015, there were on average 5,360 home health visits each year at SPCH, a three percent increase from the previous triennium. The following figure (Figure 5.9) illustrates the six year trend.

Figure 5.9: Annual home health visits, 2010 - 2015



Source: Samaritan Health Services, 2016

## Conclusion

Health insurance coverage rates have increased drastically in the past five years. In 2010, 19 percent of Lincoln County adults were uninsured, and in 2015, that rate had dropped to 4 percent. Much of the increase in insurance coverage is due to increased enrollment in the Oregon Health Plan, which grew by 6,000 people between 2010 and 2014.

The data presented in this chapter can support an initial understanding and baseline of access to medical care in Central Lincoln County, while calling attention to challenges faced by many in our community when accessing medical care.



# Chapter 6

## Morbidity and Mortality

Understanding the leading causes of illness and death is the first step on the path to preventing both the loss of life and improving the quality of life within any community. Traditional measures used to evaluate the health of populations are morbidity (incidence of disease) and mortality (deaths). Examining various cancers, heart disease, and other major causes can highlight notable improvement as well as areas in which Lincoln County is in need of improvement. The more detailed data available about disparities within particular populations and illnesses, the better communities can address these issues effectively in Lincoln County. Many of the conditions that cause illness and death within Lincoln County have well-established causes, a number of them rooted in behaviors or risk factors that can be prevented.

Many statistics are aggregated over a set of years in order to report reliable data. When incidence or prevalence rates are reported across many years, the statistic is per person per year. For example, the all-cancer incidence rate in Oregon across 2008-2012 was 448 cases per 100,000 people; this means that in each of the five years between 2008 and 2012, 448 cases were diagnosed for every 100,000 people in the population.


Morbidity and Mortality data is not available for Central Lincoln County, so Lincoln County figures are reported.

### Leading Causes of Death in Lincoln County

In 2013, the age-adjusted death rate in Lincoln County was 775 deaths per 100,000 people, compared to 717 deaths per 100,000 people in Oregon. The leading causes of death (for all ages combined) in Lincoln County are cancer, heart disease, chronic lower respiratory disease, unintentional injuries, and cerebrovascular disease (stroke). Compared with Oregon, Lincoln County has a similar number of deaths per 100,000 residents for each of its top ten causes of death (Figure 6.1).

Preventable risk factors such as tobacco use, diet, activity and alcohol use contribute substantially to these deaths. For example, in 2013, it is estimated that 30 percent of Lincoln County deaths were tobacco-related deaths. This proportion is comparable to 22 percent of tobacco-related deaths in Oregon during the same time period.<sup>128</sup>

Figure 6.1: Top 10 causes of death per 100,000 persons, age-adjusted, Lincoln County, 2013



Source: Oregon Health Authority, Oregon Public Health Assessment Tool, 2013

## Chronic Disease and Conditions

Chronic diseases, such as heart disease, stroke, cancer and diabetes are among the most prevalent, costly, and preventable of all health problems. Healthy lifestyles, such as avoiding tobacco, being physically active, and eating well, greatly reduce a person's risk for developing chronic illnesses. Research shows that access to resources that support healthy lifestyles, such as nutritious food, recreational opportunities, and high quality and affordable prevention measures (including screening and appropriate follow-up) saves lives, reduces disability, and lowers medical costs.<sup>129</sup>

### Oncology - Cancer

Cancer is the leading cause of death in Lincoln County and in Oregon.<sup>130</sup> Five types of cancer are discussed in the section: lung, colorectal, breast, prostate, and pancreatic. Lung cancer is the most common cause of cancer death for Oregonians, followed by colorectal cancer and breast cancer.<sup>131</sup> Pancreatic cancer has a very high mortality rate, in part due to the likelihood of a late diagnosis after the cancer has already progressed. Prostate cancer is a common cancer among men.

Lincoln County's annual rate of newly diagnosed cancer cases is similar to the rate in Oregon, with Lincoln County having 450 diagnoses per 100,000 individuals each year.

Lincoln County's incidence of tobacco-related cancer is significantly higher.<sup>132</sup> Data for all cancer and tobacco related cancer incidence are from different years and are therefore not directly comparable.

Cancer rates also vary between different racial and ethnic groups. In Oregon, prevalence of cancer (the proportion of the population living with cancer) varies from a low of 3.6 percent among Asians and Pacific Islanders, to a high of 11.4 percent among American Indians and Alaska Natives.

Between 2011 and 2013, the Lincoln County mortality rate from all cancers was 198 deaths per 100,000 people per year.

### **Lung and Bronchial Cancer**

Lung and bronchial cancers are closely related, and this section will combine them both as lung cancer. Lung cancer incidence in men is steadily declining as a result of decreasing smoking rates, but the incidence in women remains relatively flat.<sup>133</sup> Lung cancer is the deadliest cancer in Oregon, accounting for 27 percent of cancer deaths in the state in 2013; a number which includes tobacco and non-tobacco caused lung cancers.<sup>134</sup> The rate of lung cancer has remained fairly constant in Lincoln County and the United States over time.

In Lincoln County, smoking declined gradually from 2004 to 2011, which has major implication for cancer rates, since smoking is the leading cause of lung cancers.<sup>135</sup> The lung and bronchial cancer incidence rate in Lincoln County was 74 per 100,000 persons per year from 2008-2012. Mortality rates are also disparate in Lincoln County compared to the state. Lincoln County has a mortality rate due to lung cancer at 62 per 100,000. Oregon has a mortality rate of 47 per 100,000.

### **Breast Cancer**

Oregon has the 7<sup>th</sup> highest incidence rate for breast cancer in the United States.<sup>136</sup> Although significant improvements have occurred in early detection and treatment, breast cancer is still the leading cause of death for women in Oregon. Only a small fraction of breast cancer cases can be linked to genetics.<sup>137</sup>

The 2008-2012 age-adjusted incidence of breast cancer among women was 117 diagnoses per 100,000 women in Lincoln County compared to 128 diagnoses per 100,000 women in Oregon. In 2008-2012, the female breast cancer mortality rate in Lincoln County (27 per 100,000 women) was higher than the Oregon mortality rate due to female breast cancer (21 per 100,000 women).<sup>138</sup> Lincoln County and Oregon had mortality rates above the Healthy People 2020 target of 20.7 deaths per 100,000 females.<sup>139</sup>

## Prostate Cancer

The 2008-2012 incidence of prostate cancer in Lincoln County was 123 per 100,000 men. The mortality rate for Lincoln County, however, was among the highest in the state at 25 per 100,000 men, compared to the state mortality rate of 23 per 100,000 men.<sup>140</sup> These rates do not meet the Healthy People 2020 objective to reduce the mortality rate due to prostate cancer 22 deaths per 100,000 men.<sup>141</sup>

## Colorectal Cancer

The age-adjusted incidence of colorectal cancer in Lincoln County is 42 per 100,000 individuals, above the state incidence of 38 per 100,000 individuals. The Lincoln County mortality rate (19 per 100,000) is also above the state rate (15 per 100,000 individuals). Lincoln County has not achieved the Healthy People 2020 target to reduce the mortality rate due to colorectal cancer to 14.5 deaths per 100,000 people.<sup>142</sup>

## Pancreatic Cancer

In 2008-2012, the annual incidence rate for pancreatic cancer in Lincoln County was 12.5 per 100,000 persons.<sup>143</sup> In contrast with the other cancers discussed in this section, pancreatic cancer mortality rates are close to incidence rates, with rates of 8.9 per 100,000 in Lincoln County. Pancreatic cancer is difficult to diagnose before it has advanced, so survival rates tend to be lower than for other common cancers.

## Cancer Screening

Research shows that screening for cancer is effective in reducing serious consequences of the disease, which is generally more treatable when detected early. Breast and cervical cancer screening rates in Lincoln County is fairly consistent with state-level screening rates (Table 6.1). Additional data are needed to identify rates of screening among race/ethnic populations, age group and income level, as risk factors differ among different populations.

Table 6.1: Age-adjusted percent of cancer screening in Lincoln County and Oregon, 2010-2013

Cancer Screening Practice	Lincoln County	Oregon
<b>Mammogram within past 2 years (women 50-74 years old)</b>	77.1%	75.3%
<b>Pap test within past 3 years (women 21-65 years old)</b>	78.6%	81.7%
<b>Current on colorectal cancer screening (50-75 years old)*</b>	63.4%	61.1%

Source: Oregon Health Authority, *Health screenings among Oregon adults, 2010-2013*

\*Current on colorectal cancer screening includes the following: having a fecal occult blood test (FOBT) in the past year; a colonoscopy within the past 10 years; or, a sigmoidoscopy within the past 5 years as well as an FOBT within the past 3 years.

## Heart Disease and Stroke

After cancer, heart disease is the largest contributor to the mortality rate in Lincoln County and in Oregon. When combined with stroke and adjusted for age, diseases of the circulatory system are the leading causes of death in Lincoln County and Oregon.

### Cardiovascular Disease and Stroke

The heart attack incidence in Lincoln County is 96 per 100,000 persons.

Numerous health conditions and behaviors contribute to the potential for heart disease and stroke. The prevalence of these factors are listed below:

- High blood pressure: 35 percent,
- high blood cholesterol: 35 percent,
- diabetes: 8 percent
- obesity: 30 percent
- lack of exercise: 16 percent, and
- smoking: 33 percent.<sup>144</sup>

Many of the effects of heart disease can be reversed with healthy eating, exercise, avoidance of tobacco, and stress reduction. In addition to high blood pressure, high cholesterol, and diabetes being critical health factors of heart disease and stroke, social and economic factors are also important. For example, in the U.S., low-income adults are 50 percent more likely to suffer heart disease than top wage earners, even when other risk factors such as cholesterol or smoking, are taken into account.<sup>145</sup>

### Heart Disease Mortality

Across Oregon, the death rate for heart disease is higher in rural areas than urban areas.<sup>146</sup> Mortality rates are very different in Lincoln County; in 2013 Lincoln County (152 per 100,000 persons) had higher cardiovascular disease mortality rates than Oregon (135 per 100,000 persons).<sup>147</sup>

### Stroke Mortality

In general, stroke mortality rates in Lincoln County and in Oregon have not achieved the Healthy People 2020 target of a reduction to 34.8 deaths per 100,000 persons (Figure 6.15).<sup>148</sup> Lincoln County's mortality rate is 43 deaths per 100,000 people, as opposed to Oregon's rate of 37 deaths per 100,000 people.

## Diabetes

### Diabetes in Adults

There are two types of diabetes identified by the medical community. Type 1 diabetes is a hormonal condition in which the body does not produce enough insulin to regulate the conversion of sugar and starches into energy. Type 1 diabetes is caused by genetic and unknown factors and is usually diagnosed in children.

In Type 2 diabetes, the body develops resistance to insulin, so that dietary sugar absorbed into the bloodstream is not converted into glycogen at a healthy rate. There are both genetic risk factors and behavioral risk factors for developing type 2 diabetes. Because diabetes can cause serious health complications, it is important to prevent type 2 diabetes through healthy life choices and to catch early through health screenings.<sup>149</sup>

Prevalence of diabetes among adults in Lincoln County was 7.7 percent in 2008-2011.<sup>150</sup> This estimate may be conservative, however, as many people are unaware of their status. Diabetes often develops gradually, and symptoms and complications can take years to manifest.

### Diabetes Mortality

Overall, 2013 age-adjusted annual diabetes mortality rates have been consistently higher in Lincoln County (26 per 100,000) than they have been in Oregon (23 per 100,000). These rates, however, are lower than the national diabetes mortality rate and meet the Healthy People 2020 objective of no more than 66.6 deaths per 100,000 persons.<sup>151</sup>

## Alzheimer's Disease

Alzheimer's disease is the most common form of dementia, which is a general term for loss of memory and other intellectual abilities serious enough to interfere with daily life. Alzheimer's disease accounts for 60 to 80 percent of all cases of dementia. Alzheimer's disease is also terminal, and is the 6<sup>th</sup> most common cause of death in Lincoln County. In 2013, the Lincoln County cause-specific mortality rate of 34 deaths per 100,000 individuals was similar to the mortality rate in Oregon.

## Arthritis

Arthritis continues to be the most common cause of disability in the United States, affecting one in five Americans. Arthritis consists of over 100 different diseases and conditions that affect the joints, surrounding tissues and other connective tissues. The two most common types are osteoarthritis and rheumatoid arthritis.

The age-adjusted percentages of adults in Lincoln County who report an arthritis diagnosis is 30 percent, compared to Oregon at 25 percent.<sup>152</sup>

## Asthma

Over the past 20 years, asthma has become one of the most common chronic diseases in the United States. Oregon has one of the highest asthma rates in the nation.<sup>153</sup> Asthma results in direct health care costs (e.g., hospitalizations and emergency department visits) and indirect costs (e.g., missed school and work days and days of restricted activity), and affects the quality of life for people with asthma and their families.

### Prevalence of Asthma in Adults

For the past 10 years, the percent of Oregonians with a current asthma diagnosis has been rising slowly. Oregon ranked among the top six states for the highest percentage of adults with current asthma diagnoses in 2011.<sup>154</sup>

Two important risk factors, tobacco use and obesity, contribute to the likelihood of an asthma diagnosis. Oregon counties with asthma levels higher than the state average tend to also be counties with high smoking rates.<sup>155</sup> Likewise, counties with high levels of obesity also tend to have increased prevalence and incidence of asthma. Lincoln County's adult asthma prevalence is lower than Oregon's at 9.3 percent.<sup>156</sup>

## Infectious Diseases

Prevention and control of infectious illnesses rank among the greatest health advances of the 20<sup>th</sup> century. The World Health Organization defines infectious diseases are those that are caused by bacteria, viruses, parasites, or fungi; these diseases can be passed from person to person.<sup>157</sup> Some are transmitted via ingesting contaminated food or water. Many are spread by microorganisms in coughs or sneezes, while others result from exposures in the environment or insect bites. Diseases that spread from animals are called zoonotic infections.

All physicians, health care providers, and laboratories in Oregon are required by law to actively report confirmed or suspect diagnoses of over 50 infectious diseases and conditions to their local health departments.<sup>158</sup> These reports are directed through county health departments to the Oregon Public Health Division which collects and distributes data to inform health departments, physicians and the public. Reporting enables appropriate public health follow-up for patients, helps identify outbreaks, and provides a better understanding of disease transmission patterns. Some diseases are subject to restrictions on school attendance, day care attendance, patient care, and food handling. Communicable disease nurses in Lincoln, Benton, and Lincoln Counties investigated 1,336 reports of reportable communicable diseases during 2013, a rate of over 25 investigations every week.<sup>159</sup>

## Respiratory Illnesses

Respiratory illnesses such as the influenza virus, commonly referred to as the flu, spread from person to person when droplets from a cough or sneeze of an infected person move through the air and enter the mouth or nose of people nearby.

The common cold<sup>5</sup> and influenza are the most common respiratory illnesses. However, local, state, and national statistics for these diseases are difficult to ascertain because doctors and laboratories are not required to report them to public health authorities. This is because most people experience only mild, short-term illness, and do not seek medical attention. The illnesses are difficult to differentiate, and most are treated symptomatically rather than curatively. The Oregon Health Authority reports influenza and pneumonia mortality jointly; these rates have been steadily declining in Lincoln County, from 15 deaths per 100,000 persons in 2007 to 10 deaths in 2013.

Less common, but more serious respiratory illnesses include pneumonia, pertussis (whooping cough), and tuberculosis. In general, infectious tuberculosis is extremely rare in Lincoln County. Between 2007 and 2013, an average of 3-4 cases were reported annually.<sup>160</sup> Tuberculosis cases are actively managed and curative therapy is overseen by public health nurses.

Pertussis is a very contagious bacterial infection that causes a coughing illness which may last 6 to 10 weeks or longer. It is an endemic disease with epidemic peaks occurring every 2 to 7 years and has proven persistent despite widespread childhood immunization. There was a sharp rise of pertussis in the United States during 2012. Oregon reported more than twice as many pertussis cases in 2012 as in 2011. The number of cases of pertussis in Lincoln County fluctuates annually; an outbreak in 2012 pushed the incidence above the historical average of approximately 14 diagnoses per 100,000 people per year.

## Foodborne Illnesses

The Centers for Disease Control and Prevention (CDC) estimate that each year, 1 in 6 Americans (48 million people) get sick, 128,000 are hospitalized, and 3,000 die of foodborne diseases.<sup>161</sup> The leading causes of foodborne illness in the United States are due to exposure to norovirus, Salmonella, Campylobacter, and *Clostridium perfringens*. Norovirus, Salmonella, and Campylobacter are also among the leading causes of death due to foodborne illness.<sup>162</sup> The incidence of campylobacter in Lincoln County has historically ranged between 15 and 25 cases per 100,000 each year.

*Escherichia coli* infections, most commonly O157:H7 (a specific strain of *E. coli*), is another significant causative organism. Around 5 to 10 percent of those who are diagnosed with the

---

<sup>5</sup> More than 200 viruses cause what is typically considered the common cold, including rhinovirus, coronavirus, respiratory syncytial virus, and the parainfluenza virus.



infection develop potentially life-threatening complications. Lincoln County's rate of *E. coli* infections was 10 per 100,000 persons in 2013.<sup>163</sup>

## Sexually Transmitted Infections (STIs)

Sexually transmitted infections (STIs, also sometimes called sexually transmitted diseases, STDs) are infections that can be passed from one person to another through sexual contact. Untreated STIs can have consequences for an individual's health such as infertility and even death. Testing for STIs is a very effective mechanism for preventing the spread of STIs. Even incurable STIs, like HIV, are much less likely to be spread if those affected by the infection receive proper treatment. However, untested individuals are unable to receive the treatment they need and are also much more likely to pass on the infection to others.

Chlamydia and gonorrhea are the most common STIs in Lincoln County. Approximately 80 to 90 percent of chlamydia infections and about 50 percent of gonorrhea infections are asymptomatic in women and may go undiagnosed. If left untreated, these infections may lead to pelvic inflammatory disease, which can cause tubal infertility, ectopic pregnancy and chronic pelvic pain.<sup>164</sup>

### Chlamydia

Chlamydia is the most common reportable illness in Oregon, with infection rates steadily increasing over the past decade. In Lincoln County reported rates of chlamydia are more than twice as high in women as in men; for every 10 men diagnosed with chlamydia, 25 women are diagnosed. Current guidelines recommend chlamydia screening in women who are not symptomatic, but do not recommend the same screening for men without symptoms. This likely causes the higher rate of reported chlamydia cases among women, rather than a difference in infection rates by gender.<sup>165</sup> Overall, Lincoln County had 300 infections per 100,000 persons in 2013.

### Gonorrhea

Another reportable sexually transmitted infection that is present in Lincoln County is gonorrhea. In general, women are more likely than men to become infected with gonorrhea after exposure. However, as with chlamydia, women are less likely than men to develop symptoms following infection.<sup>166</sup> Gonorrhea infection rates in Lincoln County were 9 per 100,000 persons in 2013.

The key risk factor for sexual infections is age. Lincoln County residents between 15 and 24 years of age contract chlamydia at a rate 4.8 times higher than the infection rate among all ages. This trend holds for region infection rates as well. Gonorrhea infection rates are somewhat less influenced by age; 15-24 year olds in Lincoln County have infection rates 3.1 times as high as the infection rate among all ages (Table 6.2).

Table 6.2: Age-specific incidence rates of chlamydia and gonorrhea, diagnoses per 100,000 persons in Lincoln County, 2013

Age	Chlamydia	Gonorrhea
	Lincoln County	Lincoln County
<15	300	9
15-24	1,321	0.0
25-44	384	32
45-64	27	0.0
65+	0.0	0.0

Source: Oregon Health Authority, Oregon Public Health Assessment Tool, 2013

## HIV/AIDS

HIV/AIDS (human immunodeficiency virus/ acquired immunodeficiency syndrome) remains an important public health problem in Oregon. From 1981 through 2010, 8,753 Oregonians were diagnosed with HIV infection. Of those, 40 percent (3,540) died. Fortunately, death rates have decreased dramatically since the development of effective antiretroviral therapies. HIV/AIDS is now managed as a serious but chronic disease. As a result, the number of Oregonians living with HIV infections has increased from 2,720 in 1997 to 5,213 in 2010. New HIV diagnoses in Oregon are most common among 35–39 year old males. Between 2009 and 2013, 9 individuals were diagnosed in Lincoln County. The 5-year incidence of HIV in Lincoln County was 5 cases per 100,000 persons per year, about two-thirds of the state’s incidence (6.5 cases per 100,000 persons per year) during that time period.<sup>167</sup>

## Injury and Violence

### Violent Crime

Violent crimes are defined as offenses that involve face-to-face confrontation between the victim and the perpetrator, including homicide, rape, robbery and aggravated assault. High levels of violent crime compromise physical safety and psychological well-being. Crime rates can also deter residents from pursuing healthy behaviors such as exercising out-of-doors.<sup>168</sup>


Lincoln County had violent crime rates of 365 crimes per 100,000 people from 2010-2012. This was the second highest rate in the state. There were 132 recorded assaults in 2012 in Lincoln County.<sup>169</sup> In 2013, Lincoln County recorded 18 violent deaths, including suicide, homicide, and undetermined causes. This was a rate of 39 violent deaths per 100,000 residents, much higher than the rate in Oregon.<sup>170</sup>

## Unintentional Injury Mortality

Injuries are the number one cause of death among people under the age of 44 in Oregon and the fifth leading cause of death overall. Injury is also the number one cause of disability at all ages.<sup>171</sup> Most of the events resulting in injury, disability, or death are preventable. According to Healthy People 2020, injuries and violence have an impact on the well-being of people by contributing to premature death, disability, poor mental health, high medical costs, and high unproductivity.<sup>172</sup>

Lincoln County injury deaths follow the same pattern as the state (Figure 6.2). Falls contributed to 29 percent of accidental deaths between 2009 and 2013, followed by poisoning and motor vehicle accidents. Together, these three causes comprise 82 percent of accidental deaths in Lincoln County.

Figure 6.2: Causes of accidental injury deaths in Lincoln County, 2009-2013



Source: Oregon Public Health Assessment Tool, 2009-2013

Injury mortality is higher among males than females in all age groups in Oregon. Injury mortality rates increase with age for both sexes, starting at age 5.<sup>173</sup> The risks of different major types of injury fluctuate through a person's life. These include, among other types, falls, unintentional poisonings, motor vehicle accidents, and self-harm.

## Mental Health Conditions

Mental health disorders are experienced by people of all ages, from early childhood through old age. Research suggests that only about 17 percent of U.S. adults are considered to be in a state of optimal mental health. An estimated 26 percent of Americans age 18 years and older are living with a mental health disorder in any given year, and 46 percent will have a mental health

disorder during their lifetime.<sup>174</sup> These disorders include, among others, anxiety, depression, behavior disorders, persistent suicidal thoughts, schizophrenia, and Alzheimer’s disease.<sup>175</sup> County Health Rankings reports the number of poor mental health days each month, both as a proxy for mental health diagnoses and as an indicator of overall mental wellness. Residents of Lincoln County reported an average of 3.6 poor mental health days over the previous month. This measure is based on survey responses to the question: “Thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?”<sup>176</sup>

Oregonians across the state reported an average of 3.3 poor mental health days. The Healthy People 2020 benchmark is 2.3, placing Lincoln County and the state in the worst 10 percent for this measure, with clear room for improvement. From 2008 to 2011, 60 percent of Lincoln County residents reported no poor mental health in the past 30 days. These rates are statistically equivalent to the statewide rate of 65 percent.<sup>177</sup> From 2010 to 2013, self-reported depression rates in Lincoln County were 28 percent. Lincoln County depression rates were close to the state rate of 25 percent.<sup>178</sup>

## Suicide

Suicide is a death resulting from the intentional use of force against oneself. As a public health concern, it relates to both injury and violence, and mental health. However, while many unintentional injuries can be prevented by making one’s environment safer, suicide can also be effectively prevented by providing treatment to those with mental health disorders. Therefore, suicide is discussed in the context of mental health. Suicide is an important public health problem in Lincoln County. It is also the leading cause of injury-related death in the state and is the 9th leading cause of death for Oregonians. There are more deaths in Oregon due to suicide than due to car crashes. The suicide rate in Lincoln County was 30 suicides per 100,000 residents. The statewide rate in 2013 was 18 per 100,000 persons.<sup>179</sup> Additional detail is given in Chapter 7.

## Conclusion

Understanding the leading causes of illness and death is a first step on the path to preventing both the loss of life and improving the quality of life within Lincoln County. While leading causes of death in Lincoln County closely mirror those of the state, examining various cancers, heart disease, and other major causes reveal areas of vast improvement, as well as areas in which Lincoln County is doing more poorly than the state average. Data on many sub-populations are noticeably absent throughout this chapter. As discussed throughout the chapter, many of the conditions that cause illness and death within Lincoln County have well-established causes, with a number of them rooted in behaviors or risk factors that can be prevented. The following chapter takes a closer look at behaviors and risk factors that affect a person’s health and well-being across the life course.

# Chapter 7

## Health Across the Life Course

A life course framework helps to illuminate the ways in which experiences during key stages of life contribute to health outcomes throughout an individual's lifetime.<sup>180</sup> This framework builds upon the previously discussed social determinants of health to illustrate that a person's environment and the systems in which they live can affect health outcomes differently during different stages of life. For example, there are ways in which maternal and infant disparities contribute to childhood and adolescent experiences which, in turn, contribute to adult and older adult health outcomes. This can have effects not only on an individual's life, but can also span generations, creating and contributing to persistent disparities within the community. By taking a look at the ways in which certain life stages and health factors interact, new opportunities to improve community health can be uncovered. This chapter addresses health behaviors such as engaging in physical activity; maintaining healthy eating habits; being tobacco-free; and using alcohol and prescription drugs appropriately. This larger view illuminates ways that people protect and promote health for others, including assuring a healthy start for children, preventing and managing chronic conditions, preventing disease and injury, and promoting good mental health.

This chapter is organized to follow the course of a person's life. Factors that influence the health of a mother have a lasting effect on the health of her children as infants, adolescents, and adults. The behavioral and lifestyle choices of children and adolescents affect their wellbeing throughout life, as do other health factors that arise during childhood, such as oral care and mental health. Adults continue to influence their health by adopting healthy behaviors or discontinuing unhealthy ones. They may now feel the effects of health conditions that began earlier in life but took time to develop, or reap the benefit of the healthy choices that they have made. Living a healthy life as one ages is very possible, and it is never too late to improve one's health. However, the elderly are also at risk for health issues that may not affect younger people, including falls, mental decline, or elder abuse.

Central Lincoln County data is not available for these data, so Lincoln County figures are reported.


## Maternal and Infant Health

Since healthy aging starts at the beginning of one’s life, public health professionals can assess the health of a community by starting with the health of mothers and infants during and immediately after pregnancy. This section takes a closer look at maternal and infant health outcomes and various factors that impact them. All fertility data is based on the county of residence, not the county where the infant was born.

### Fertility Rate (Total Fertility Rate, TFR)

The total fertility rate (TFR) is the total number of births per 1,000 women in a given year. The TFR is based on the age-specific fertility rates of women in their “child-bearing years”, which is ages 15 to 44. Lincoln County has a TFR that is higher than the state’s. Among racial/ethnic groups, women who identify as Hispanic or Latina have the highest TFR in Lincoln County of 100 births per 1,000 women (Figure 7.1).


Figure 7.1: Fertility rate, total (births per 1,000 women) by race/ethnicity in Lincoln County, 2011-2013



Source: Oregon Healthy Authority, Center for Health Statistics, Birth Certificate Data, 2011-2013

Note: Fertility rate data is based on county of residence, not county of birth

Figure 7.2: Age-specific fertility rates (births per 1,000 women) by maternal age in Lincoln County, 2011-2013




Source: Oregon Health Authority, Center for Health Statistics, Birth Certificate Data, 2011-2013

## Prenatal Care and Healthy Pregnancy

In Lincoln County from 2011 to 2013, a total of 86 percent of all mothers were able to access adequate prenatal care. As shown in Figure 7.3, younger mothers are less likely to access adequate prenatal care than older mothers. Compared with women between 25 and 39, women under the age of 25 are nearly twice as likely to receive inadequate or no prenatal health care in Lincoln County.<sup>181</sup>


Figure 7.3: Percent of mothers accessing inadequate or no prenatal care in Lincoln County by age group, 2011-2013



Source: Oregon Health Authority, Center for Health Statistics, Birth Certificate Data, 2011-2013

There also exist disparities in prenatal care access among mothers of different race/ethnic groups in Lincoln County. Overall, mothers who identify as White, non-Hispanic tend to access adequate prenatal care more frequently when compared to all other racial/ethnic groups (Figure 7.4).<sup>182</sup>

Figure 7.4: Percent of mothers accessing inadequate or no prenatal care by race/ethnicity in Lincoln County, 2011-2013



Source: Oregon Health Authority, Center for Health Statistics, Birth Certificate Data, 2011-2013

## Unhealthy Behavior during Pregnancy

### Smoking during Pregnancy

Smoking during pregnancy is the single most preventable cause of illness and death among infants. Smoking during pregnancy increases the risk of stillbirth, low birth weight, sudden infant death syndrome (SIDS), and preterm birth. It also contributes to cognitive and behavioral problems, and respiratory problems in both the mother and the child.<sup>183</sup>

On average in 2011-2013, 19 percent of mothers smoked during pregnancy in Lincoln County (Table 7.1). However, there is a notable difference in smoking rates when comparing age groups, in which the rate of smoking among pregnant women under the age of 25 is over two times the rate of smoking among pregnant women over the age of 25.



Table 7.1: Maternal smoking rates (percentages) among pregnant women by age in Lincoln County, 2011-2013

Population (by age)	Lincoln County
15 to 17	16 %
18 to 19	22 %
20 to 24	30 %
25 to 29	18 %
30 to 34	13 %
35 to 39	10 %
40 to 44	12 %
<b>Total</b>	<b>19 %</b>

Source: Oregon Health Authority, Center for Vital Statistics, 2011-2013

Smoking cessation counseling and programs offered during prenatal care can provide effective assistance to encourage pregnant women to quit smoking. There currently are no established smoking cessation programs specifically for mothers in Lincoln County, but efforts are being made to make them available at the county level.

### Alcohol Use During Pregnancy

Drinking alcohol during pregnancy can cause miscarriage, stillbirth, and a range of lifelong disorders, known as fetal alcohol spectrum disorders (FASDs). Children with FASDs can have a host of problems, including poor coordination, hyperactivity behavior, difficulty paying attention, poor memory, difficulty in school, learning disabilities, speech and language delays, poor reasoning and judgment skills, vision or hearing problems, and complications with the heart, kidney, or bones. There is no known safe amount of alcohol to drink during pregnancy, and no known safe time to drink alcohol during pregnancy.<sup>184</sup>


The Pregnancy Risk Assessment Monitoring System (PRAMS), a national surveillance system, provides information about women who have had a recent live birth. Oregon state-level data indicates that 92 percent of pregnant mothers abstained from alcohol during the last 3 months of their pregnancies. Less than one percent had more than one drink per week during the third trimester.<sup>185</sup> There are no Lincoln County data available at present.

### Teen Parenting

Teen mothers are less likely to receive early prenatal care, and are more likely to experience blood-pressure complications and premature birth.<sup>186</sup> Children of teenage mothers are also more likely to become teen parents themselves, be incarcerated during adolescence, drop out of school, experience more health problems, and are two times as likely to experience abuse and neglect. Negative effects of early childbearing on teenage fathers include an increased likelihood of partaking in delinquent behaviors, such as alcohol and drug abuse or dealing, and fewer years of completed school in comparison to their childless peers.<sup>187</sup>


The most recent information available suggests that, overall, Lincoln County teen pregnancy rates (ages 15 to 17 and 18 to 19) have decreased between 2009 and 2013 (Figures 7.6 and 7.7). Given the small number of teen pregnancies each year, three year averages are shown. The three year average in 2008-2010 among 15-17 year-olds was 26 pregnancies per 1,000 women age 15-17. This number declined to 22.5 pregnancies per 1000 women age 15-17 in 2011-2013. This decline is similar among 18-19 year old women; the rate declined from 109 pregnancies per 1,000 women age 18-19 in 2008-2010 to 72 pregnancies per 1,000 women age 18-19 in 2011-2013. Lincoln County teen pregnancy rates were above state teen pregnancy rates.

Figure 7.6: Pregnancy rate per 1,000 women age 15-17 years in Lincoln County, 3 year moving average, 2008-2013



Source: Oregon Health Authority, Center for Health Statistics, Birth Certificate Data, 2008-2013

Figure 7.7: Pregnancy rate per 1,000 women age 18-19 years in Lincoln County, 3 year moving average, 2008-2013




Source: Oregon Health Authority, Center for Health Statistics, Birth Certificate Data, 2008-2013

Disparities in teen pregnancy rates emerge when the overall Lincoln County figure is broken down. For example, despite the overall decline in rates, there are striking differences in teen birth rates for Hispanic and non-Hispanic populations. Between 2011 and 2013, Hispanic teens aged 15 to 19 had a pregnancy rate in Lincoln County that was eight percent lower than that of non-Hispanic teens (Figure 7.8). Notwithstanding the greater Hispanic teen pregnancy rates in

Lincoln County, the pregnancy rate among Hispanic teens are declining faster than the pregnancy rate among non-Hispanic teens.

Figure 7.8: Pregnancy rate, Hispanic versus non-Hispanic, women age 15-19 years in Lincoln County, 3 year moving average, 2008-2013



Source: Oregon Health Authority, Center for Health Statistics, Birth Certificate Data, 2008-2013

## Infant Mortality

The annual infant mortality<sup>6</sup> occurrence in Lincoln County was 3 deaths per 1,000 births from 2011 to 2013. Infant mortality rates are lower in Lincoln County (3 per 1,000 births) than in Oregon (4.98 per 1,000 births).<sup>188</sup> Lincoln County has surpassed the Healthy People target of 6.0 per 1,000 births.<sup>189</sup>

## Premature Birth and Low Birth Weight

Premature birth, and low birth weight among infants are commonly used measures of maternal and infant health. Infants that are born too early and/or with a low birth weight are at higher risk of dying in the first year of life and of having developmental problems and worse health outcomes throughout life.<sup>190,191</sup> Both conditions are preventable to varying degrees and have been found to be influenced by a variety of factors.

<sup>6</sup> Infant mortality is defined as the death of a live-born infant before the age of 1.

## Premature Birth

Premature birth (also known as preterm birth) is a measure of births that occur before the projected full term of the pregnancy. Infants are considered premature when they are born before completing 37 weeks (about 8.5 months) of pregnancy.<sup>192</sup>

In Lincoln County, the percent of preterm births is 8.5 percent, which is well below the Healthy People 2020 target of 11.4 percent.<sup>193</sup> However, disparities exist among women when stratified by race/ethnicity, as shown below in Table 7.2.

Table 7.2: Percent of births that are premature in Lincoln County, 2011-2013

Lincoln County	
<b>White, non-Hispanic</b>	7.9%
<b>Black/African American, non-Hispanic</b>	20%*
<b>American Indian/Alaska Native</b>	1.6%*
<b>Asian &amp; Pacific Islander</b>	16.7%*
<b>Hispanic/Latino</b>	11.5%
<b>All births</b>	8.5%

Source: Oregon Health Authority, Center for Health Statistics, Birth Certificate Data, 2011-2013

\* This number may be statistically unreliable due to small numbers and should be interpreted with caution

## Low Birth Weight

Low birth weight results when an infant fails to grow sufficiently during pregnancy and can both signal and cause health problems with the infant. Infants are considered to have low birth weight if they weigh less than 2,500 grams (about 5.5 pounds at birth).

From 2011 to 2013, approximately 7.9 percent of all infants born in Lincoln County had a low birth weight, which almost meets the Healthy People 2020 target of 7.8 percent.<sup>194</sup> Differences exist among racial/ethnic groups within the county. Table 7.3 illustrates the variation across different racial/ethnic groups within Lincoln County.

Table 7.3: Percent of infants born with low birth weight by race/ethnicity in Lincoln County, 2011-2013

Lincoln County	
<b>White, NH</b>	6.7%
<b>Black/African American, NH</b>	20%*
<b>American Indian/Native American, NH</b>	4.9 %*
<b>Asian &amp; Pacific Islander, NH</b>	16.7%*
<b>Hispanic/Latino</b>	11.5%
<b>All infants</b>	7.9%

Source: Oregon Health Authority, Center for Health Statistics, Birth Certificate Data, 2011-2013

\* This statistic may be statistically unreliable due to small numbers and should be interpreted with caution

## Breastfeeding

Breastfeeding is associated with numerous health benefits for infants, such as boosting immune system response, reducing the risk of Type 2 diabetes, and preventing obesity. Breastfeeding also promotes maternal-child bonding. The American Academy of Pediatrics recommends exclusively breastfeeding for the first 6 months after birth and further recommends continued breastfeeding for a year or more after birth.<sup>195</sup>

### Breastfeeding in Lincoln County

Data on breastfeeding are limited at both the state and county level. However, state programs, such as the Nutrition and Health Screening Program for Women, Infants, and Children (WIC), give some insight into the percentage of participating women who breastfeed. Table 7.4 displays the available county data on mothers who participate in the WIC program and the rate of breastfeeding.<sup>196</sup>

Table 7.4: Breastfeeding rates among WIC mothers in Lincoln County, 2014

	Lincoln County
<b>Percent of pregnant women served by WIC</b>	71%
<b>Percent of WIC mothers who started out breastfeeding (initiation)</b>	90%
<b>Percent of WIC mothers who breastfed exclusively for 6 months</b>	21%

Source: Oregon Health Authority, 2014 WIC Facts

In addition to WIC, most health care providers encourage women to breastfeed their children, and there are many breastfeeding classes and support groups available in Lincoln County.

## Childhood and Adolescence

Childhood and adolescence are formative times in a person's life. The number and severity of adverse experiences during childhood affects an individual's risk for alcoholism, depression, heart disease, liver disease, intimate partner violence, sexually transmitted infections, smoking, and suicide. Adverse events include emotional, physical, and sexual abuse and neglect, and various types of household dysfunctions such as violence against mothers, substance abuse, mental illness, parental separation or divorce, or an incarcerated household member.<sup>197,198</sup>

## Gender Identity and Sexual Orientation


Adolescence is a time of developing sexual awareness and gender expression, although many children are aware of their developing gender identity from a very early age. Because most state and national surveys do not ask questions related to sexual orientation or gender identity, it is difficult to estimate the health needs of lesbian, gay, bisexual, transgender, or queer (LGBTQ) children, youth and adults in Lincoln County and Oregon.

Available data include survey responses on harassment among adolescents in our public schools. In Lincoln County during the 2013-2014 school year, 8<sup>th</sup> and 11<sup>th</sup> graders reported having been harassed by a peer who thought they were gay, lesbian, bisexual, or transgender more frequently than 11<sup>th</sup> graders. Overall, harassment based on perceptions about sexual orientation declines with age.<sup>199,200, 201</sup>

## Child Abuse

In 2014, there were a total of 348 reports of child abuse/neglect in Lincoln County, of which 118 (20 percent) were founded (determined to be abuse). There were an additional 141 unfounded reports, and 45 indeterminable reports.<sup>202</sup> The types of abuse/neglect include mental injury, physical/medical neglect, physical abuse, sexual abuse, sexual exploitation, or threat of harm. Most often, the perpetrators of child abuse and neglect are family members (93.9 percent of reports); parents account for 78.2 percent of all perpetrators.<sup>203</sup> In recent years, the rate of child abuse in Lincoln County has decreased and is similar to the rate in Oregon.<sup>204,205,206,207</sup>

Figure 7.10: Founded abuse rate per 1,000 for children under 18 years of age in Lincoln County, 2010-2014




Source: Oregon Department of Human Services, Child Welfare Data Book 2011, 2012, 2013, 2014

Rates include neglect, physical abuse, and sexual abuse

\*2012 data is from the Portland State University Population Research Center. Starting in 2013, the population data is one year behind the year shown and is from Puzanhera, C., Sladky, A. and Kang, W. (2014). "Easy Access to Juvenile Populations: 1990-2013."

Not all reported cases of child abuse result in a foster care placement. Children are placed in foster care for a variety of reasons. Some are placed in foster care because their families cannot provide them with basic safety and protection, while others have had negative experiences such as parental substance abuse, sexual or physical abuse, and abandonment. In Oregon, many children are in foster care due to a history of abuse or neglect.<sup>208</sup>

Figure 7.11: Children in foster care, rate per 1,000 children in Lincoln County, 2011-2014



Source: Oregon Department of Human Services: Children, Adults and Families Division (2012). *Child Welfare Data Book*

\*State totals do not include Title IV-E eligible children served by tribes.


## Bullying/Peer Abuse

Violence in schools can affect the learning environment and contribute to absenteeism. Students who are bullied, harassed, and feel unsafe or otherwise victimized, are more likely to miss classes, skip school, feel depressed or exhibit problem behaviors. Research shows that comprehensive discipline, positive behavioral support and anti-bullying programs in schools can reduce the incidence of harassment among primary and secondary school students.<sup>209</sup>

In 2015, ten percent of 8<sup>th</sup> graders and four percent of 11<sup>th</sup> graders did not go to school at least once in the past 30 days due to feeling unsafe at school or on their way to school.<sup>210</sup>

Figure 7.12 below shows that reasons for harassment at school differ among age groups at the county level, and that the overall incidence of harassment among Lincoln County students is common. While the percent of students who report having been harassed at school in the past month tends to decrease with age, reasons for and severity of harassment vary among age groups. Aside from all or other reasons, harassment for physical characteristics is the most reported reason for harassment across all age groups.<sup>211</sup>

Figure 7.12: Percent of students in 8<sup>th</sup> and 11<sup>th</sup> grade, who experienced bullying in the past 30 days by reason in Lincoln County, 2015



Source: Oregon Healthy Teens Survey, 2015

## Psychological Distress

Mental health includes our emotional, psychological, and social well-being, and is essential to the overall health and wellbeing of an individual. The World Health Organization defines it as “a state of well-being in which the individual realizes her or her own abilities, can cope with the normal stresses of life, can work productively and fruitfully, and is able to make a contribution to his or her community.”<sup>212</sup>

In Oregon, suicide rates are higher than the national average and about 70 percent of people who died by suicide from 2003 to 2012 also had depression. Among all age groups, the suicide rate in 2013 per 100,000 people was 19 in Lincoln County.<sup>213</sup>

The following table highlights the percentage of 8<sup>th</sup> and 11<sup>th</sup> grade students in Lincoln County and Oregon that exhibited signs of depression, thought about suicide, or attempted suicide during 2015 (Table 7.5). The rate of attempted suicide is higher among 8<sup>th</sup> graders in Lincoln County than among 11<sup>th</sup> graders in Lincoln County.<sup>214</sup>



Table 7.5: Percent of 8<sup>th</sup> and 11<sup>th</sup> grade students that exhibited signs of depression, thoughts about suicide, or actually attempted suicide during the last 12 months, Lincoln, and Oregon, 2015

	Grade	Lincoln	Oregon
<b>Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities</b>	8th	36 %	27 %
	11th	28 %	29 %
<b>Seriously considered attempting suicide</b>	8th	19 %	16 %
	11th	16 %	16 %
<b>Attempted suicide at least once</b>	8th	15 %	8 %
	11th	5 %	6 %

Source: Oregon Healthy Teens Survey, 2015

## Physical Activity

Regular physical activity helps improve overall health and wellness, reduces risk for obesity, and lessens the likelihood of developing many chronic diseases including diabetes, cancer, and heart disease. National physical activity guidelines recommend that children engage in at least 60 minutes of physical activity each day, including aerobic, muscle strengthening, and bone strengthening activity.

The Healthy People 2020 objective for physical activity aims to increase the proportion of adolescents who meet current national physical activity guidelines to 32 percent.<sup>215</sup> 21 percent of 8<sup>th</sup> graders and 33 percent of 11<sup>th</sup> graders met national guidelines.<sup>216</sup>

Table 7.6 shows that the majority of youth in 8<sup>th</sup> and 11<sup>th</sup> grade in Lincoln County do not spend more than two hours per school day watching television. Among 11<sup>th</sup> graders, a larger percentage of youth in Lincoln County spend less than two hours watching television than Oregon youth overall. Lincoln County surpasses the HP 2020 target for this measure for youth in 11<sup>th</sup> grade.<sup>217</sup>

Table 7.6: Percent of youth who view television for no more than two hours per school day in Lincoln County and Oregon, 2015

Grade	Lincoln County	Oregon
<b>8<sup>th</sup> Grade</b>	73 %	76 %
<b>11<sup>th</sup> Grade</b>	80 %	80 %

Source: Oregon Healthy Teens Survey, 2015

Table 7.7 shows that about one-half to two-thirds of 8<sup>th</sup> and 11<sup>th</sup> graders in Lincoln County spend less than 2 hours per day on the computer or on their phone. These rates are somewhat

below to the state average, and fall well short of the Healthy People 2020 target of 82.6 percent.<sup>218</sup>

Table 7.7: Percent of youth who play video/computer games or use a computer for something that is not school work for no more than two hours per school day (including time spent on social networks and on smartphones) in Lincoln County and Oregon, 2015

Grade	Lincoln County	Oregon
<b>8<sup>th</sup> Grade</b>	42 %	54 %
<b>11<sup>th</sup> Grade</b>	55 %	58 %

Source: Oregon Healthy Teens Survey, 2015

## Nutrition

There is a well-established link between eating a healthy and balanced diet, and an increasing number of health benefits. A healthy and balanced diet involves eating a variety of foods which provide essential nutrients (like dietary fiber and potassium), in the right amount – with negative health consequences from consuming too little or too much food.<sup>219</sup> In addition to promoting health, and supporting a healthy weight, mounting evidence links a healthy diet to lowered risks of chronic disease, including several types of cancer, osteoporosis, and cardiovascular disease.<sup>220</sup>

Table 7.8: Percent of youth consuming at least 5 servings of fruits and vegetables per day in Lincoln County and Oregon, 2015

Grade	Lincoln County	Oregon
<b>8<sup>th</sup> Grade</b>	23 %	23 %
<b>11<sup>th</sup> Grade</b>	22 %	20 %

Source: Oregon Healthy Teens Survey, 2015

## Obesity

Being obese or overweight<sup>7</sup> is a complicated health condition. The risk of unhealthy weight is influenced by diet, exercise, and other behaviors, but it also depends strongly on genetic and environmental factors. Obesity is also correlated with socio-economic status and other social determinants of health. In addition to being a poor health outcome, obesity and overweight status can increase the risk of many diseases such as diabetes, heart disease, and possible cancer.

<sup>7</sup> Obesity is defined as having a body mass index (BMI) of 30 or more; Overweight is defined as having a BMI of above 25 and less than 30. Healthy weight is a BMI between 20 and 25.

The Oregon Healthy Teens Survey<sup>8</sup> found that nearly forty percent of all eighth graders in Lincoln County are overweight or obese (Table 7.9). Lincoln County youths are generally more likely to be overweight than Oregon youths, with the exception of 11<sup>th</sup> graders in Lincoln County. The rate of being overweight does not change much between 8<sup>th</sup> and 11<sup>th</sup> grade, but obesity rates decrease by almost half.<sup>221</sup>

Table 7.9: Overweight and obesity prevalence in Lincoln County and Oregon, 2015

	Lincoln	Oregon
<b>8<sup>th</sup> grade overweight</b>	20 %	15 %
<b>8<sup>th</sup> grade obese</b>	19 %	11 %
<b>11<sup>th</sup> grade overweight</b>	19 %	15 %
<b>11<sup>th</sup> grade obese</b>	10 %	13 %

Source: Oregon Healthy Teens Survey, 2015

## Oral Health

Another childhood health issue that has ramifications for the rest of a person’s life is adequate dental care. Good oral health is essential to overall physical and mental health, and encompasses more than just dental check-ups. Oral disease can lead to cavities and gum ailments, which can in turn contribute to other diseases or conditions. Conversely, certain chronic mental and physical health conditions can also contribute to declines in oral health. Poor oral health can also affect self-esteem, reduce employment opportunities, and increase absenteeism.<sup>222</sup>

Cavities are almost completely preventable through optimal water fluoridation, application of dental sealants to children’s teeth, effective oral hygiene (brushing teeth and flossing), and regular preventive visits to the dentist.<sup>223</sup> The proportion of 8<sup>th</sup> grade and 11<sup>th</sup> grade youth who have ever had a cavity is higher than the Healthy People 2020 target of no more than 48.3 percent (Table 7.10). The proportions do not change much in the three years between 8<sup>th</sup> grade and 11<sup>th</sup> grade – this indicates that most tooth decay occurs in children before the 8<sup>th</sup> grade.<sup>224</sup>

Table 7.10: Percent of youth who have ever had a cavity in Lincoln County and Oregon, 2015

Grade	Lincoln County	Oregon
<b>8<sup>th</sup> Grade</b>	75 %	69 %
<b>11<sup>th</sup> Grade</b>	77 %	75 %

Source: Oregon Healthy Teens Survey, 2015

<sup>8</sup> The Oregon Healthy Teens Survey distributes a questionnaire to 8<sup>th</sup> and 11<sup>th</sup> graders; therefore, adolescent data is richest for these age groups.

Achieving and maintaining good oral health is a significant challenge for many people in Lincoln County, particularly those with lower incomes. This challenge may be exacerbated by the fact that no cities, districts, or water supplies in Lincoln County are fluoridated (see Chapter 3: Environment).

One of the objectives of Healthy People 2020 is to increase the proportion of U.S. communities with fluoridated water to 75 percent.<sup>225</sup> Approximately 27 percent of Oregon residents have access to fluoridation through community water systems, the second lowest percentage in the country.<sup>226</sup> However, no municipal or public water systems in Lincoln County are fluoridated.<sup>227</sup>

## Alcohol, Tobacco, and Prescription and Illicit Drug Abuse

Alcohol and prescription medications are consumed appropriately and responsibly by most of the population. However problems frequently occur when these substances are over-consumed, used inappropriately, combined with other substances, or consumed while engaging in risky activities such as driving or unsafe sexual activity. The costs to society of the misuse of alcohol, prescription medications and other drugs are massive, and include injury and death due to overdose; effects on unborn children of drug users; impacts on family, crime and homelessness; spread of infectious disease including through sexual transmission and needle sharing; and financial costs associated with lost productivity, healthcare, and legal expenses for individuals and the wider community.<sup>228</sup>

Table 7.11: Percent of youth who reported taking part in illicit activities in the past 30 days in Lincoln County and Oregon, 2015

	Grade	Lincoln	Oregon
<b>Consumed at least one alcoholic beverage</b>	8th	17 %	12 %
	11th	24 %	30 %
<b>Consumed at least 5 alcoholic beverages within a couple of hours</b>	8th	6 %	5 %
	11th	15 %	17 %
<b>Smoked cigarettes</b>	8th	8 %	4 %
	11th	7 %	9 %
<b>Used e-cigarettes</b>	8th	10 %	9 %
	11th	10 %	17 %
<b>Used marijuana</b>	8th	22 %	9 %
	11th	23 %	19 %
<b>Used prescription drugs without a doctor's orders</b>	8th	15 %	4 %
	11th	15 %	7 %

Source: Oregon Healthy Teens Survey, 2015

## Alcohol Use

The younger a person begins drinking regularly, the greater the chance that person will develop a clinically defined alcohol disorder. Youth who start drinking before the age 15, compared to those who start at 21, are far more likely to be injured while under the influence of alcohol, to be in a motor vehicle crash after drinking, or to become involved in a physical fight after drinking.<sup>229</sup> Overall, alcohol use among Lincoln County youth tends to increase with age, reflecting the state trend displayed in Table 7.11.

## Binge Drinking


Middle and high school youth in Lincoln County and Oregon report binge drinking at similar rates. Approximately 6 percent of Lincoln County 8<sup>th</sup> graders reported binge drinking in 2015 (Table 7.11, above). This rate increased to 15 percent among 11<sup>th</sup> graders.<sup>230</sup> Lincoln County likely meets the Healthy People 2020 objective of reducing the percent of high school seniors (12<sup>th</sup> graders) who binge drink to below 23 percent, but it is not possible to directly compare the rates between 11<sup>th</sup> graders and 12<sup>th</sup> graders.<sup>231</sup>

## Tobacco Use among Adolescents

Tobacco products are designed to deliver nicotine, an addictive drug that changes the way the brain works, causing tobacco users to crave repeated doses. Youth are especially sensitive to nicotine and can become dependent more quickly than adults. Because of their dependency, nearly three out of four teen smokers continue using tobacco products into adulthood.<sup>232</sup>

Tobacco use is present among Lincoln County youth but has been decreasing over time. Figure 7.13 below illustrates this trend, in addition to showing that smoking among youth increases with age.<sup>233,234</sup>

Figure 7.13: Percent of students that reported having smoked cigarettes at least once in the past 30 days, Lincoln County, 2007 and 2015



Source: Oregon Healthy Teens Survey, 2007 and 2015

2015 was the first year the Oregon Healthy Teens Survey asked students about electronic cigarette use. The rate of electronic cigarette use among 8<sup>th</sup> graders was higher than that of cigarette smoking (10 percent versus 8 percent), as was the rate among 11<sup>th</sup> graders (10 percent versus 7 percent).

## **Marijuana, Prescription Drug, and Illicit Drug Use**

Recreational marijuana is still illegal for all individuals under 21 years of age. The effects of marijuana on children and adults have not been studied to the degree that other legal substances have been, including alcohol and cigarettes. Another major public health concern is the abuse of prescription drugs. When these drugs are misused or taken without a doctor's prescription they can be just as harmful as illegal street drugs.

Among youths in Lincoln County, marijuana use was generally more than twice as prevalent as cigarette smoking as shown in Table 7.11. Marijuana use did not increase from 8<sup>th</sup> grade to 11<sup>th</sup> grade. In Lincoln County, one out of every five 11<sup>th</sup> graders surveyed reported using marijuana in the past 30 days. Adolescents in Lincoln County abuse prescription drugs at rates higher than the state.<sup>235</sup> There are no reliable data on other illicit drug use among adolescents in the county.

Many healthy behaviors that children learn continue to support their health as adults, including good oral hygiene, exercise and nutritious diets, and abstaining from drugs or tobacco. The next section discusses the prevalence of these behaviors among adults in Lincoln County.

## **Adults**

As individuals enter adulthood, they become independent in their decision making. With adulthood comes a new set of stressors and risks, such as the potential for partner violence or easier access to alcohol, tobacco, and drugs. At the same time, working adults have more independence and resources to engage in healthy behaviors such as consuming nutritious food and getting adequate exercise.

Many of the topics that were covered in the childhood health section are revisited here, and some additional areas are introduced. Much of the data reported in this section covers adults from age 18 onward.

## **Domestic Violence**

Domestic violence, which includes many forms of abuse, affects children and adults. Physical abuse, sexual abuse or assault, intimidation, verbal abuse and emotional abuse, or threats of such harm are all forms of domestic violence. Domestic violence can include abuse from a

household member (including roommates or caregivers), intimate partners (including dating partners) or a family member (whether or not they live with the victim).<sup>236</sup>

My Sister's Place (MSP)<sup>237</sup> is a non-profit organization serving residents of Lincoln County that provides supportive services to victims of domestic violence, sexual assault, and dating abuse. Services include crisis intervention, emergency shelter, 24-hour crisis line, safety planning, advocacy, court information and support, agency and resource referrals, education, peer counseling, and outreach activities.

My Sister's Place (MSP)<sup>238</sup> is a non-profit organization serving residents of Lincoln County that provides supportive services to victims of domestic violence, sexual assault, and dating abuse. Services include crisis intervention, emergency shelter, 24-hour crisis line, safety planning, advocacy, court information and support, agency and resource referrals, education, peer counseling, and outreach activities.

Domestic violence not only has an effect on the victim, but can also have an effect on children; domestic violence poses a threat to children's emotional, psychological, and physical well-being. Children who live with domestic violence are also at an increased risk to become direct victims of child abuse.<sup>239</sup>

## **Abuse of Vulnerable Adults**

Vulnerable adults include the elderly and adults of all ages with physical or mental disabilities, whether living at home or being cared for in a health facility. Abuse and maltreatment of vulnerable adults can include physical, emotional, or sexual abuse, caregiver neglect, and financial exploitation. The information in this section includes adults and seniors.

In Lincoln County there were 4 allegations of abuse against adults with an intellectual and/or developmental disability in 2014.


## **Mental and Emotional Health**

Mental and emotional health begins in childhood and extends throughout a person's life. Many mental disorders manifest during the transition from adolescence to adulthood, including depression, schizophrenia, and bipolar disorder.

A major depressive episode (MDE) is defined as a period of at least two weeks when a person experienced a depressed mood or a loss of interest or pleasure in daily activities. In addition to this, four of the seven symptoms reflecting the criteria for major depressive disorder (as described in the 4<sup>th</sup> edition of the DSM-IV) must have been experienced by the individual.<sup>240</sup> From 2008 to 2010, eight percent of Lincoln County adults age 18 years and older reported having had a MDE in the past year.<sup>241,242</sup> This is comparable to seven percent of Oregon adults aged 18 years and older having reported an MDE in the past year.<sup>243</sup>

The most serious consequence of poor mental health is suicide. Overall, the suicide rate among Oregon adult males is 3.7 times the rate among adult females. This is a larger ratio than in Lincoln County, where the suicide rate among men is 3.1 times the rate among women.<sup>244</sup> The total suicide rate increases with age, but this is due primarily to the outsize effect of male suicide rates, which increase with age. Among males of all age groups in Lincoln County from 2009 to 2012, males over the age of 65 had the highest suicide rate at 58 per 100,000 men (Figure 7.14). After the age of 70, the rate of suicide among older adult men rises dramatically (up to 72 per 100,000 men for males 85 and older in the state of Oregon).<sup>245</sup> Females had a much lower rate of suicide, averaging 9 per 100,000 women, and this rate does not increase with increasing age. The suicide rate among women peaks at 15 per 100,000 women between the ages of 45 and 64.<sup>246</sup>

Figure 7.14: Suicide rates per 100,000 men, per 100,000 women, and per 100,000 individuals, Lincoln County age adjusted averages, 2009-2013



Source: Oregon Health Authority, Oregon Public Health Assessment Tool, 2009-2013



## Physical activity

Physical activity is important for maintaining health as a person ages. Recommendations for adults include at least an hour and fifteen minutes of vigorous-intensity activity or two-and-a-half hours of moderate-intensity activity every week, in addition to muscle-strengthening activities on two or more days a week.<sup>247</sup>

Overall, 18 percent of adults in Lincoln County met the CDC guidelines for physical activity<sup>9</sup> from 2010-2013, compared to 25 percent of adults in Oregon.<sup>248</sup>

Of greater concern is the proportion of the population that gets little to no exercise. In Lincoln County, 16 percent of adults report no physical activity outside of work, compared to the Oregon average of 18 percent.<sup>249</sup>

At the state level, participation in physical activity varies by race/ethnicity, household income, and by level of education. Adults with less than a high school education, those earning less than \$24,999, and Latinos are less likely to meet CDC physical activity recommendations than their peers.<sup>250</sup> As with children and youth, county-level data that describe physical activity levels among adults by race/ethnicity or level of household income are not available.

Nutrition and eating habits are frequently set early in life. Good nutrition can delay the physical signs of aging and prevent or slow the development of many chronic diseases, including diabetes and cancer. Approximately one in four adults in Lincoln County and in Oregon consumes at least 5 servings of fruits and vegetables per day.<sup>251</sup>

## Obesity

Obesity is a complicated health issue, with many factors that affect the likelihood of obesity, and many downstream health issues that are influenced by body weight.

While about 30 percent of children in Lincoln County are overweight or obese, the prevalence of overweight or obesity more than doubles among adults. An estimated 25 percent of adults in Lincoln County are obese; an additional 30 percent are overweight.<sup>252</sup> Therefore, about 65 percent of Lincoln County adults are either overweight or obese.

## Alcohol, Tobacco, and Drug Use

Data shows that younger adults (particularly in their 20's) are at a higher risk for misuse of alcohol and other drugs. One particular area for concern is the misuse of prescription drugs. Misuse of these drugs is highest among young adults (aged 18 to 25).<sup>253</sup> As the most commonly abused type of prescription drugs, painkillers provide a useful marker for prescription drug

---

<sup>9</sup> The CDC recommends 30 minutes of moderate physical activity on five or more days per week.

misuse trends. While data shows little change in the self-reported pain experienced by Americans, the amount of painkillers dispensed in the U.S. has quadrupled since 1999, as have the deaths resulting from prescription painkillers.

Of particular concern is the rate of misuse of prescription painkillers among Medicaid patients. Research shows higher rates of inappropriate provider prescribing practices and patient use as compared to privately insured patients. Potential inappropriate prescribing practices include overlapping prescriptions for painkillers, high daily doses, or long-acting painkillers, and have been shown in at least one study to be present in particularly high rates amongst Medicaid enrollees.<sup>254</sup>

### Alcohol Use

Excessive drinking is a risk factor for many adverse health outcomes, such as hypertension, alcohol poisoning, unintended pregnancy, fetal alcohol syndrome, inter-personal violence, and motor vehicle crashes.<sup>255</sup> It can also contribute to a number of health issues including heart disease and stroke, high blood pressure, cirrhosis, coma, and even death.<sup>256</sup> The following data includes adults ages 18 and older.

Excessive drinking is defined differently for men and women, due to different metabolic rates and average body weights. Among men, excessive drinking is defined as 2 or more alcoholic drinks per day for a period of 30 days. In Lincoln County, 14 percent of men reported excessive drinking (Table 7.12).

For women, excessive drinking is defined as one or more alcoholic drinks per day for a period of 30 days. In general, excessive drinking among women is lower in Lincoln County than among men (Table 7.12).

Table 7.12: Alcohol abuse among adults, 18 years and older in Lincoln County and Oregon, 2010-2013

	Gender	Lincoln	Oregon
<b>Consumed at least two alcoholic beverages per day for the past 30 days</b>	Male	14 %	6 %
<b>Consumed at least one alcoholic beverage per day for the past 30 days</b>	Female	6 %	7 %
<b>Consumed at least 5 (male) or 4 (female) alcoholic beverages within a couple of hours in the past 30 days</b>		16 %	18 %

Source: Oregon BRFSS 2010-2013

### Binge Drinking among Adults


For adults over the age of 18, binge drinking is defined as consuming five or more drinks at one time for men and four or more drinks at one time for women.<sup>257</sup> Binge drinking is more common across Lincoln County and in the state than drinking every day. Sixteen percent of Lincoln County adults reported binge drinking within the past month in 2013 (Table 7.13).<sup>258</sup>

Lincoln County meets the Healthy People 2020 objective of reducing the percent of adults that reported having engaged in binge drinking within the past month to 24.4 percent.<sup>259</sup>

## Tobacco Use and Exposure to Secondhand Smoke

Tobacco use is the single most preventable cause of disease, disability, and death in the United States. Tobacco use in any form can cause serious diseases and health problems, including cancers of the lung, bladder, kidney, pancreas, mouth, and throat, heart disease and stroke, lung diseases (i.e., emphysema, bronchitis, and chronic obstructive pulmonary disease), pregnancy complications, gum disease and vision problems.<sup>260</sup>

Figure 7.15: Age-adjusted percent of adults who currently smoke cigarettes Lincoln County, 2002-2011



Source: Oregon BRFSS, Table II: Prevalence of Modifiable Risk Factors among Adults, 2002-2011

More recent data has been collected that includes up to 2013, but should not be compared to the data in Figure 7.15 above due to changes in survey methodology. The 2010-2013 data is presented in Table 7.13 below, and is likely more accurate due to a more representative survey sample. These data indicate that Lincoln County is not meeting the Healthy People 2020 smoking target of 12 percent or below.

Table 7.13: Age-adjusted percent of adults who currently smoke cigarettes in Lincoln County and Oregon, 2010-2013

	Lincoln County	Oregon
<b>Smoking rates</b>	33% %	19 %

Source: Oregon BRFSS, Table II: Prevalence of Modifiable Risk Factors among Adults, 2010-2013

## Secondhand Smoke Exposure

Secondhand smoke is a mixture of the smoke exhaled by a person smoking, and the smoke from burning tobacco in a cigarette, pipe, or cigar. Secondhand smoke contains the same toxic chemicals and carcinogens as inhaled tobacco smoke, and even brief exposure has been found to put a nonsmoker's health at risk. In adults, secondhand smoke exposure has been found to cause lung cancer and heart disease. Children exposed to secondhand smoke are more at risk for ear infections, asthma attacks, respiratory symptoms and infections, and a greater risk for sudden infant death syndrome (SIDS).<sup>261</sup> Measures to reduce the amount of secondhand smoke exposure to others include, but are not limited to, quitting smoking, forbidding smoking in the home, and forbidding smoking in a shared car. Table 7.14 illustrates that the majority of households in Lincoln County have rules against smoking in the home, while a smaller majority hold the same rules against smoking in family cars.

Table 7.14: Family tobacco/secondhand smoke practices (age-adjusted) in Lincoln County and Oregon, 2008-2011

	Lincoln County	Oregon
<b>Have rules against smoking in the home</b>	91 %	91 %
<b>Have rules against smoking in family cars</b>	70 %	82 %

Source: Oregon Health Authority, Table IV: Age-adjusted and unadjusted prevalence of tobacco use among adults, 2008-2011

## Older Adult Health


As people age, many of the behaviors they have adopted over their lives begin to pay dividends, whether it is exercising regularly, eating healthy food, or avoiding drugs and tobacco. Older adults also encounter a different set of challenges for staying healthy. This section revisits some of the ongoing behaviors and also highlights new factors that affect health.

### Preventing Falls

Falls are a major cause of injury and hospitalization, and the 10<sup>th</sup> leading cause of death among older Oregonians.<sup>262</sup> Nearly one in three older adults experiences a fall each year, and 20-30 percent of those who fall suffer injuries. As commonly as they occur, injuries and deaths due to falls are not an inevitable consequence of aging; they can be prevented. Muscle weakness is a significant contributing factor in falls, so physical activity is widely viewed as among the most important interventions for preventing injuries related to falls among older adults.

Hospitalization rates for falls increase drastically as adults age; the rate of hospitalizations due to a fall for adults 75 years and older is more than six times the rate for adults 60-74 years. Older adults hospitalized for falls are nearly six times more likely to be discharged into long term care compared to older adults hospitalized for other conditions. In 2013, the cost for fall injury hospitalization among adults 65 years and older in Oregon totaled to more than \$219 million.<sup>263</sup> Between 2011 and 2013, the mortality rate from falls in Lincoln County was 79 deaths per 100,000 residents age 65 and older. Figure 7.16 below highlights the difference in mortality rates for different age groups among the elderly in Lincoln County.

Figure 7.16: Fall mortality among elder adults in Lincoln County, 2011-2013



Source: Oregon Health Authority, Oregon Public Health Assessment Tool, 2011-2013

### Depression and suicide

Lincoln County suicide rates are also higher among older adults, with 27 suicides per 100,000 adults age 65 and older between 2009 and 2013. This rate is marginally higher than the age adjusted rate for all Lincoln County residents. This increased rate conceals the difference between older men and women, however. The suicide rate among older men was 5 percent higher than among all men and was lower than the suicide rate among men age 45-64. The suicide rate among older women was twice as high as among all women, and equal to the 45-64 age rate. See Figure 7.14 in the Adults Mental and Emotional Health section for a visual representation of these data.

### Conclusion

There were a number of promising changes in behavioral indicators over the past triennium. Prenatal smoking rates declined from 23 percent to 15 percent, 11<sup>th</sup> grade smoking rates declined from 22 percent to 7 percent, and smoking rates continued to decline among all adults. However, 10 percent of 11 graders reported using e-cigarettes, in the first year that

these data were recorded. Binge drinking among 11<sup>th</sup> graders increased by about 6 percent over the previous triennium. Behavioral factors among adults did not change much, other than a sharp decrease in the percent of adults getting the CDC recommended level of physical activity. However, this change was mostly due a higher level of activity recommended by the CDC. The other major change was the apparent rise in smoking rates among adults; however, it is likely that this increase (from 27 percent to 33 percent) is due to a change in survey methodology. The previous triennium may have underreported smoking rates.

The life course framework helps us recognize two key components of community health: the health issues which are most important at particular stages of life, and when they are best addressed. At the same time, we are able to identify particular disparities that are present throughout the life course, such as the far-reaching risks of poor health tied to low birth weight. In this context, differences in low birth weight that exist by race in Lincoln County, as well as differences in related behaviors, like smoking among pregnant women, become particularly important for health improvement efforts.

As we gather more data on the health behaviors and practices discussed in this chapter, we will be able to make distinctions between such factors as race/ethnicity, age, and income status. Such data would enable us to focus prevention efforts at the age and sub-population level at which they are most needed in Lincoln County, resulting in countless health benefits throughout the life course.

# Conclusion

## Meeting Challenges with Resources

As highlighted throughout the Community Health Needs Assessment, there are many factors that influence and affect health outcomes, both positively and negatively, in Central Lincoln County. The CHNA provides an opportunity to identify the many health concerns, disparities and impacts that residents face in their daily lives.

A health assessment is truly important to help identify needs and opportunities for improvement. At the same time, it is important to highlight the various resources and assets that are alive and well within our communities. These resources and assets refer to the many types of human, social and economic resources that Central Lincoln County can offer to address problems. Organizations, agencies and partners within and across the three counties can collaborate to improve the health and quality of life for residents.

### General Health Status

In 2016, Lincoln County was ranked 29 out of 36 counties for health outcomes, and 24 out of 36 for health factors.<sup>264</sup> It is clear in these numbers that Lincoln County has a lot of opportunity ahead to work on improving overall health status for the residents who live here. The County Health Rankings look at the different factors and conditions that affect the health and well-being of county residents, and are made up of four categories: health behavior, clinical care, social and economic factors, and physical environment.

In the 2016 County Health Rankings report, Lincoln County had strengths in its primary care physician ratio, the proportion of residents who had attended college, and clean air.<sup>265</sup>

### Knowledge and Skills in Caring for and Promoting Health

Central Lincoln County is part of a three-county region that shares a long history of collaboration and partnership among various organizations and agencies to improve and promote health.

- Samaritan Pacific Communities Hospital and Samaritan Health Services work tirelessly to improve the health of the people of Central Lincoln County by providing excellent health care and supporting social programs.
- Lincoln County's Tobacco Prevention & Education Program aims to reduce tobacco-related illness and death. There also exist other population-based prevention and chronic disease programs that reduce the onset and incidence of many chronic conditions and help residents take control of their health.

- Lincoln County is home to a variety of medical care, dental care, vision care, elder care, medical clinics, doctors, nurse practitioners, and alternative medicine which can be expanded upon to meet the needs of all residents.

## Social Support Networks

- Lincoln County has a comprehensive network of social support and opportunity for the aging population.
- The county offers specialized support for people with mental illness, addictions, disabilities, and children with behavioral or emotional problems.
- The county has a strong commitment to the health and wellbeing of children and youth. This commitment includes a focus on issues such as increasing family stability, kindergarten readiness, and equitable service coordination. Numerous organizations exist to address education, nutrition, and social support for children and families.
- Community Services Consortium serves as the community action agency supporting the most vulnerable populations in the county.

Without being able to call out every organization and project that supports the health of the Central Lincoln County, what is shown above only highlights a few examples; each example is the result of efforts by countless community partners. A wealth of collective action and resources exists within Lincoln County. Overcoming the many health challenges facing residents depends on this collective action and the vitally important part that each of our community partners play.

## Resources

- Lincoln County has an excellent basic framework to assist homeless persons (i.e. emergency shelter, transitional housing, and permanent affordable housing).
- The county is particularly strong in offering a wide choice in public schools, private schools, and alternative schooling opportunities.
- The county has several service providers which provide adult education (i.e. literacy, GED and parenting courses).
- The county is particularly strong in offering job seeking services, vocational training, and general support for unemployed persons.
- The county maintains safe, well-marked roads and bike lanes that help prevent traffic injuries and chronic disease.
- The county has a history of collaboration among various sectors to promote many successful and progressive transportation and built environment programs (i.e. Alternative mode options, Dial-A-Bus, PDX transit, Safety sidewalk and ramp program, Public Transit).



## Evaluation Impact


Samaritan Pacific Communities Hospital has provided services and supports to address the health needs prioritized in the previous community health needs assessment completed in 2013. Through both internal and external activities focused on the nine community benefit areas, SPCH utilizes a process evaluation method to determine community impact. The most significant health needs identified in the prior community health needs assessment included obesity, access to health care, substance abuse, homelessness, mental health, literacy, alcohol abuse and tobacco use.

A variety of actions were taken to address the identified needs, such as the offering health fairs, workshops and classes to the community that addressed obesity, literacy, homelessness, substance abuse and tobacco use. Since 2013 over 16,000 residents participated in one or more of the aforementioned events. To address the access to health care, SPCH has provided transportation, interpretative services, increased medical office hours, expanded urgent care site hours and offered screenings, exams and complete physicals in local schools, after-school programs and in Senior Centers. During this same timeframe over 600 residents received interpretative services during a medical appointment and over 2,000 were screened or examined. More detailed information on activities that support the priorities are contained in the 2013, 2014 and 2015 Community Benefit Plans. SPCH also provided over \$300,000 to non-profit organization through direct financial support or in-kind contributions.

## Community priorities

Samaritan Pacific Communities Hospital reached out to community members through a series of key informant interviews, focus groups, and surveys to learn about the health priorities of the community. Full results are presented in the appendix to this document. A summary of the survey results is shown on the following page. Community members were asked to rate 26 health priorities on a scale of 1 to 5, where 1 is lowest priority and 5 is highest priority. The results were averaged for each of the 26 categories.

Figure 8.13: Community Health Prioritizations



A total of **633** respondents from Lincoln County answered the question: “For the following issues that affect health, please circle how much attention you think they should get in our communities on a scale of 1 to 5” from the Community Health Perceptions Survey. Responses are reported as mean values. Of those that responded, the top five issues were identified as: “Access to mental health care” with a mean value of **4.6**; “Access to medical care” with a mean value of **4.3**; “Poverty” with a mean value of **4.3**; “Child abuse or neglect” with a mean value of **4.3**; and lastly “Housing costs” with a mean value of **4.3**”.

- 
- <sup>1</sup> World Health Organization (WHO). (1985). Constitution of the World Health Organization. Retrieved from <http://www.who.int/about/definition/en/print.html>
  - <sup>2</sup> Oregon Health Authority, Office for Oregon Health Policy and Research. (2011). Research Brief: Health Equity. Retrieved from <http://www.oregon.gov/oha/oei/docs/health-equity-brief.pdf>
  - <sup>3</sup> Oregon Health Authority, Public Health Division. (2010). The Burden of Asthma in Oregon: 2010. Retrieved from [http://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Asthma/Documents/burden/or\\_asthma2010.pdf](http://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Asthma/Documents/burden/or_asthma2010.pdf)
  - <sup>4</sup> Oregon Health Authority, Public Health Division. (2012). Oregon overweight, obesity, physical activity and nutrition facts. Retrieved from [http://public.health.oregon.gov/PreventionWellness/PhysicalActivity/Documents/Oregon\\_PANfactst\\_2012.pdf](http://public.health.oregon.gov/PreventionWellness/PhysicalActivity/Documents/Oregon_PANfactst_2012.pdf)
  - <sup>5</sup> Hillemeier, M., Lynch, J., Harper, S., & Casper, M. (2004). Data Set Directory of Social Determinants of Health at the Local Level [data set]. Retrieved from [http://www.cdc.gov/dhdsdp/docs/data\\_set\\_directory.pdf](http://www.cdc.gov/dhdsdp/docs/data_set_directory.pdf)
  - <sup>6</sup> King County. (2012). Equity and social justice annual report. Retrieved from [www.kingcounty.gov/equity](http://www.kingcounty.gov/equity)
  - <sup>7</sup> Minnesota Department of Health and Healthy Minnesota Partnership. (2012). The Health of Minnesota: Statewide Health Assessment. Retrieved from [www.health.state.mn.us/healthymnpartnership](http://www.health.state.mn.us/healthymnpartnership)
  - <sup>8</sup> Public Broadcasting Service. (2008). Unnatural Causes: is inequality making us sick? Retrieved from [http://www.unnaturalcauses.org/assets/uploads/file/AmazingFacts\\_small.pdf](http://www.unnaturalcauses.org/assets/uploads/file/AmazingFacts_small.pdf)
  - <sup>9</sup> Oregon Health Authority. (2013). 10-Year Plan for Oregon Project: Healthy People Policy Vision. Retrieved from <http://www.oregon.gov/COO/Ten/docs/PeopleOutcome.pdf>
  - <sup>10</sup> US Census Bureau (2011) State and County Quick Facts: Lincoln County, OR. Retrieved from <http://quickfacts.census.gov/qfd/states/41/41043.html>
  - <sup>11</sup> U.S. Census Bureau. (2013). Selected Social Characteristics in the United States, American Community Survey 3-Year estimates, 2011-2013, Table DP02. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_DP02&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_DP02&prodType=table)
  - <sup>12</sup> U.S. Census Bureau. (2013). Selected Social Characteristics in the United States, American Community Survey 3-Year estimates, 2011-2013, Table DP02. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_DP02&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_DP02&prodType=table)
  - <sup>13</sup> U.S. Census Bureau. (2013). Selected Social Characteristics in the United States, American Community Survey 3-Year estimates, 2011-2013, Table DP02. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_DP02&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_DP02&prodType=table)
  - <sup>14</sup> U.S. Census Bureau. (2013). Veteran Status, American Community Survey 5-year estimates, 2009-2013, Table S2101. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_5YR\\_S2101&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_5YR_S2101&prodType=table)
  - <sup>15</sup> U.S. Census Quick Facts. (2011). State & County QuickFacts: Veterans. Retrieved from [http://quickfacts.census.gov/qfd/meta/long\\_VET605211.htm](http://quickfacts.census.gov/qfd/meta/long_VET605211.htm)
  - <sup>16</sup> U.S. Census Bureau. (2013). Selected Social Characteristics in the United States, American Community Survey 3-Year estimates, 2011-2013, Table DP02. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_DP02&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_DP02&prodType=table)
  - <sup>17</sup> U.S. Census Bureau. (2013). Selected Social Characteristics in the United States, American Community Survey 3-Year estimates, 2011-2013, Table DP02. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_DP02&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_DP02&prodType=table)
  - <sup>18</sup> U.S. Census Bureau. (2000). Total Population, Universe: Total population, Census 2000 Summary File 1 (SF 1) 100-percent data, Table P001. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=DEC\\_10\\_SF1\\_P1&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=DEC_10_SF1_P1&prodType=table)
  - <sup>19</sup> Centers for Disease Control and Prevention, Minority Health. (2012). Definitions. Retrieved from <http://www.cdc.gov/minorityhealth/populations/REMP/definitions.html#Def>
  - <sup>20</sup> U.S. Census Bureau. (2013). Selected Social Characteristics in the United States, American Community Survey 3-Year estimates, 2011-2013, Table DP02. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_DP02&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_DP02&prodType=table)
  - <sup>21</sup> Pew Research Latino Center (2014). Lincoln County, Oregon. Retrieved from <http://www.pewhispanic.org/states/county/41043/>
  - <sup>22</sup> Pew Research Latino Center (2014). Benton County, Oregon. Retrieved from <http://www.pewhispanic.org/states/county/41003/>
  - <sup>23</sup> Pew Research Latino Center (2014). Lincoln County, Oregon. Retrieved from <http://www.pewhispanic.org/states/county/41041/>
  - <sup>24</sup> Oregon Department of Education. (n.d.). Student Ethnicity statistics, academic year 2013-2014. Retrieved from <http://www.ode.state.or.us/sfda/reports/r0067Select.asp>
  - <sup>25</sup> U.S. Equal Employment Opportunity Commission, U.S. Department of Justice Civil Rights Division. (2008). Americans with Disabilities Act: Questions and Answers. Retrieved from <http://www.ada.gov/qandaeng.htm>
  - <sup>26</sup> Watson, A., Hanrahan, P., Luchins, D., Lurigio, A. (2001). Mental Health Courts and the Complex Issue of Mentally Ill Offenders. *Psychiatric Services*, 52(4), 477-481. Retrieved from <http://dx.doi.org/10.1176/appi.ps.52.4.477>
  - <sup>27</sup> U.S. Census Bureau. (2013). Disability Characteristics, American Community Survey 5-Year estimates, 2009-2013, Table S1810. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_5YR\\_S1810&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_5YR_S1810&prodType=table)
  - <sup>28</sup> U.S. Census Bureau. (2006). American Community Survey Content Test Report: Evaluation Report Covering Disability. Retrieved from [https://www.census.gov/content/dam/Census/library/working-papers/2007/acs/2007\\_Brault\\_01.pdf](https://www.census.gov/content/dam/Census/library/working-papers/2007/acs/2007_Brault_01.pdf)

- 
- <sup>29</sup> U.S. Census Bureau. (2013). ACS Demographic and Housing Estimates, American Community Survey 3-Year estimates, 2011-2013, Table DP05. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_DP05&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_DP05&prodType=table)
- <sup>30</sup> World Health Organization. (2014). Ambient (outdoor) air quality and health. Retrieved from <http://www.who.int/mediacentre/factsheets/fs313/en/>
- <sup>31</sup> National Association of County & City Health Officials (NACCHO). (2013). Statement of Policy: Healthy Food Access. Retrieved from <http://www.naccho.org/advocacy/positions/upload/13-04-Healthy-Food-Access-2.pdf>
- <sup>32</sup> National Oceanic and Atmospheric Administration. (2015). Climate Data Online. Retrieved from: <http://www.ncdc.noaa.gov/cdo-web/>
- <sup>33</sup> National Oceanic and Atmospheric Administration. (2015). Climate Data Online. Retrieved from: <http://www.ncdc.noaa.gov/cdo-web/>
- <sup>34</sup> United States Department of Agriculture. (2015). Natural resources conservation service SNOTEL data. Retrieved from [http://www.nrcs.usda.gov/wps/portal/nrcs/detail/or/snow/products/?cid=nrcs142p2\\_046165](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/or/snow/products/?cid=nrcs142p2_046165)
- <sup>35</sup> Oregon Climate Service. (2015). Oregon Climate Data. Retrieved from <http://www.ocs.orst.edu/oregon-climate-data>
- <sup>36</sup> Oregon Secretary of State. (2015). Oregon Beach Bill Records. Retrieved from <http://sos.oregon.gov/archives/Pages/records/landmark-beach.aspx>
- <sup>37</sup> Active Living Research. (2011). Research Brief: The Power of Trails for Promoting Physical Activity in Communities. Retrieved from [http://activelivingresearch.org/files/ALR\\_Brief\\_PowerofTrails\\_0.pdf](http://activelivingresearch.org/files/ALR_Brief_PowerofTrails_0.pdf)
- <sup>38</sup> City of Newport Bicycle and Pedestrian Advisory Committee. (2008). Newport Pedestrian and Bicycle Plan. Retrieved from [http://www.thecityofnewport.net/dept/pln/documents/Newport-Pedestrian-Bicycle-Plan\\_July2008.pdf](http://www.thecityofnewport.net/dept/pln/documents/Newport-Pedestrian-Bicycle-Plan_July2008.pdf)
- <sup>39</sup> Centers for Disease Control and Prevent. (2015). Community Health Status Indicators: Lincoln County, OR. Retrieved from <http://wwwn.cdc.gov/CommunityHealth/profile/currentprofile/OR/Lincoln/>
- <sup>40</sup> Centers for Disease Control and Prevent. (2015). Community Health Status Indicators: Benton County, OR. Retrieved from <http://wwwn.cdc.gov/CommunityHealth/profile/currentprofile/OR/Benton/>
- <sup>41</sup> Centers for Disease Control and Prevent. (2015). Community Health Status Indicators: Lincoln County, OR. Retrieved from <http://wwwn.cdc.gov/CommunityHealth/profile/currentprofile/OR/Lincoln/>
- <sup>42</sup> Community Prevention Services Task Force. (2013). Preventing Dental Caries: Community Water Fluoridation: Task Force Finding and Rationale Statement. Retrieved from <http://www.thecommunityguide.org/oral/supportingmaterials/RRfluoridation.html>
- <sup>43</sup> Centers for Disease Control and Prevention. (1999). Achievements in Public Health, 1900-1999: Fluoridation of Drinking Water to Prevent Dental Caries. *MMWR*, 48(41), 933-940. Retrieved from <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm4841a1.htm>
- <sup>44</sup> Center for Disease Control. (2012). Community Water Fluoridation, 2012 Water Fluoridation Statistics. Retrieved from <http://www.cdc.gov/fluoridation/statistics/2012stats.htm>
- <sup>45</sup> Centers for Disease Control and Prevention. (2013). Designing and Building Healthy Places. Retrieved from <http://www.cdc.gov/healthypaces/>
- <sup>46</sup> Centers for Disease Control and Prevention. (2013). Indoor Environmental Quality. Retrieved from <http://www.cdc.gov/niosh/topics/indoorenv/>
- <sup>47</sup> Berkeley Lab, Indoor Air Quality Scientific Findings Resource Bank. (2013). Health Risk of Dampness or Mold in Houses. Retrieved from <http://www.iaqscience.lbl.gov/dampness-risks-house.html>
- <sup>48</sup> Personal communication with Benton County TPEP coordinator. (2015). Tobacco Prevention and Education Program Plan Update.
- <sup>49</sup> Personal communication with Lincoln County TPEP coordinator. (2015). Tobacco Prevention and Education Program Plan Update.
- <sup>50</sup> Lincoln County, Oregon. (2013). Smoking at County Facilities. Lincoln County Ordinances 7.550 – 7.580. Retrieved from [http://www.co.lincoln.or.us/sites/default/files/fileattachments/county\\_counsel/page/384/2013-lcc-chapter-07.pdf](http://www.co.lincoln.or.us/sites/default/files/fileattachments/county_counsel/page/384/2013-lcc-chapter-07.pdf)
- <sup>51</sup> Personal communication with Benton County TPP coordinator. (2015). Tobacco Prevention and Education Program Plan Update.
- <sup>52</sup> Personal communication with Lincoln County TPEP coordinator. (2015). Tobacco Prevention and Education Program Plan Update.
- <sup>53</sup> Lincoln County, Oregon. (2013). Smoking at County Facilities. Lincoln County Ordinances 7.550 – 7.580. Retrieved from [http://www.co.lincoln.or.us/sites/default/files/fileattachments/county\\_counsel/page/384/2013-lcc-chapter-07.pdf](http://www.co.lincoln.or.us/sites/default/files/fileattachments/county_counsel/page/384/2013-lcc-chapter-07.pdf)
- <sup>54</sup> Robert Wood Johnson Foundation, County Health Rankings. (2015). Retrieved from <http://www.countyhealthrankings.org/>
- <sup>55</sup> Oregon Health Authority. (2015). Environmental Public Health Tracking. Retrieved from <https://public.health.oregon.gov/HealthyEnvironments/TrackingAssessment/EnvironmentalPublicHealthTracking/Pages/index.aspx>
- <sup>56</sup> Oregon Food Bank. (2011). Rural Grocery Store Owner and Customer Assessment: Benton County. Retrieved from [http://www.oregonfoodbank.org/~/\\_media/files/community-food-systems/rural%20grocer%20reportfinaloptpdf.pdf](http://www.oregonfoodbank.org/~/_media/files/community-food-systems/rural%20grocer%20reportfinaloptpdf.pdf)
- <sup>57</sup> Robert Wood Johnson Foundation, County Health Rankings. (2015). Retrieved from <http://www.countyhealthrankings.org/>
- <sup>58</sup> Oregon Health Authority. (2015). Environmental Public Health Tracking. Retrieved from <https://public.health.oregon.gov/HealthyEnvironments/TrackingAssessment/EnvironmentalPublicHealthTracking/Pages/index.aspx>
- <sup>59</sup> [http://www.who.int/social\\_determinants/en/](http://www.who.int/social_determinants/en/)
- <sup>60</sup> World Health Organization. (2011). Rio Political Declaration on Social Determinants of Health. *Geneva: WHO*. Endorsed by WHO Member States at the Sixty-fifth World Health Assembly (WHA) in Geneva, Switzerland in May 2012. Retrieved from <http://www.who.int/sdhconference/declaration/en/>.
- <sup>61</sup> <http://www.improvingpopulationhealth.org/blog/2012/04/the-link-between-income-and-health.html>
- <sup>62</sup> <http://www.nber.org/reporter/spring03/health.html>
- <sup>63</sup> Institute for Policy Studies. (2015). Inequality and Health. Retrieved from <http://inequality.org/inequality-health/>

- <sup>64</sup> <https://www.census.gov/hhes/www/poverty/about/overview/measure.html>
- <sup>65</sup> US Census Bureau (2009-2013). Table S1701 Poverty Status in the Past 12 months: American Community Survey 3-Year Estimates. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_S1701&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_S1701&prodType=table)
- <sup>66</sup> Centers for Disease Control and Prevention: Morbidity and Mortality Weekly Report. (2013). CDC Health Disparities and Inequalities Report – United States, 2013. Retrieved from <http://www.cdc.gov/mmwr/pdf/other/su6203.pdf>
- <sup>67</sup> US Census Bureau (2009-2013). Table S1701 Poverty Status in the Past 12 months: American Community Survey 3-Year Estimates. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_S1701&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_S1701&prodType=table)
- <sup>68</sup> U.S. Census Bureau. (2013). Table S1701: Poverty Status in the Last 12 Months, 2011-2013 American Community Survey 3-Year Estimates.
- <sup>69</sup> U.S. Department of Health and Human Services: Office of the Assistant Secretary for Planning and Evaluation. (2015). 2015 Poverty Guidelines. Retrieved from <http://aspe.hhs.gov/2015-poverty-guidelines>
- <sup>70</sup> <http://www.commissiononhealth.org/PDF/Oe8ca13d-6fb8-451d-bac8-7d15343aacff/Issue%20Brief%204%20Dec%2008%20-%20Work%20and%20Health.pdf>
- <sup>71</sup> State of Oregon Employment Department. (2015). Economic Data: Unemployment Rates: Local Area Unemployment Statistics (LAUS). Retrieved from <https://www.qualityinfo.org/ed-uesti/?at=1&t1=4101000000,4121010540,4121018700,4104000041~unemprate~y~2005~2015>
- <sup>72</sup> State of Oregon Employment Department. (2015). April 2015 Employment and Unemployment in Oregon's Counties. Retrieved from <https://www.qualityinfo.org/documents/10182/73818/Labor+Force+and+Unemployment+by+Area?version=1.10>
- <sup>73</sup> U.S. Census Bureau. (2013). Means of Transportation to Work by Industry, American Community Survey 3-Year Estimates, 2011-2013, Table B08126. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_B08126&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_B08126&prodType=table)
- <sup>74</sup> [http://www.npc.umich.edu/publications/policy\\_briefs/brief9/](http://www.npc.umich.edu/publications/policy_briefs/brief9/)
- <sup>75</sup> [http://www.bls.gov/emp/ep\\_chart\\_001.htm](http://www.bls.gov/emp/ep_chart_001.htm)
- <sup>76</sup> Braveman, P., Sadegh-Nobari, T., & Egarter, S. (2011). Issue brief series: Exploring the social determinants of health: Early childhood experiences and health. Retrieved from [eric.ed.gov](http://eric.ed.gov)
- <sup>77</sup> Benefits.gov. (2015). Oregon Head Start. Retrieved from <http://www.benefits.gov/benefits/benefit-details/1935>
- <sup>78</sup> Oregon Department of Education, Early Learning Division. (2014). Oregon Head Start Prekindergarten Programs 2013-2014 Directory. Retrieved from <http://www.ode.state.or.us/superintendent/priorities/2013-14-opkhsdirectory---final.pdf>
- <sup>79</sup> Oregon State University. (2012). Child Care and Education in Lincoln County. Retrieved from <http://health.oregonstate.edu/sites/default/files/occrp/pdf/Lincoln-County-Child-Care-Profile-2012.pdf>
- <sup>80</sup> Oregon State University. (2012). Child Care and Education in Benton County. Retrieved from <http://health.oregonstate.edu/sites/default/files/occrp/pdf/Benton-County-Child-Care-Profile-2012.pdf>
- <sup>81</sup> Oregon State University. (2012). Child Care and Education in Lincoln County. Retrieved from <http://health.oregonstate.edu/sites/default/files/occrp/pdf/Lincoln-County-Child-Care-Profile-2012.pdf>
- <sup>82</sup> McCarty, C.A., Mason, W.A., Kosterman, R., Hawkins, J.D., Lengua, L.J., & McCauley, E. (2008). Adolescent school failure predicts later depression among girls. *Journal of Adolescent Health*, 43(2), 180-187.; Bogart, L.M., Collins, R.L., Ellickson, P.L., and Klein, D.J. Are adolescent substance users less satisfied with life as young adults and if so, why? *Social Indicators Research*, 81(1), 149-169
- <sup>83</sup> Robert Wood Johnson Foundation, Commission to Build a Healthier America. (2011). Issue Brief 5: Exploring the Social Determinants of Health: Education and Health. Retrieved from [http://www.rwjf.org/content/dam/farm/reports/issue\\_briefs/2011/rwjf70447](http://www.rwjf.org/content/dam/farm/reports/issue_briefs/2011/rwjf70447)
- <sup>84</sup> Oregon Department of Human Services, Oregon Tobacco Prevention and Education Program. (2007). Oregon adults who have lower income or have not finished high school data report-2007. Retrieved from [www.public.health.oregon.gov/PreventionWellness/TobaccoPrevention/Documents/lowincomefact.pdf](http://www.public.health.oregon.gov/PreventionWellness/TobaccoPrevention/Documents/lowincomefact.pdf)
- <sup>85</sup> U.S. Census Bureau. (2013). Table S1501: Educational Attainment, 2011-2013. American Community Survey 3-Year Estimates.
- <sup>86</sup> Anderson, S.A. (1990). Core indicators of nutritional state for difficult to sample populations. *The Journal of Nutrition*, 120(11), 1555-1600.
- <sup>87</sup> Bhattacharya, J., Currie, J., & Haider, S. (2004). Poverty, food insecurity, and nutritional outcomes in children and adults. *Journal of Health Economics*, 23, 839-862.
- <sup>88</sup> Department of Agriculture: Food and Nutrition Service. (2013). Child Nutrition Programs; Income Eligibility Guidelines. Retrieved from [http://www.fns.usda.gov/sites/default/files/IEG\\_Table-032913.pdf](http://www.fns.usda.gov/sites/default/files/IEG_Table-032913.pdf)
- <sup>89</sup> U.S. Department of Health and Human Services: Office of the Assistant Secretary for Planning and Evaluation. (2015). 2015 Poverty Guidelines. Retrieved from <http://aspe.hhs.gov/2015-poverty-guidelines>
- <sup>90</sup> Feeding America. (2010). Map the Meal Gap, Food Insecurity in Your County: Lincoln County, Oregon. Retrieved from [www.feedingamerica.org/mapthegap](http://www.feedingamerica.org/mapthegap)
- <sup>91</sup> Oregon Health Authority. (2015). WIC Eligibility Guidelines. Retrieved from <https://public.health.oregon.gov/HealthyPeopleFamilies/wic/Pages/income.aspx>
- <sup>92</sup> Oregon Health Authority. (2014). 2014 WIC Facts: Lincoln County Health Department. Retrieved from [https://public.health.oregon.gov/HealthyPeopleFamilies/wic/Documents/annual/annual\\_Lincoln.pdf](https://public.health.oregon.gov/HealthyPeopleFamilies/wic/Documents/annual/annual_Lincoln.pdf)
- <sup>93</sup> Oregon Health Authority. (2014). 2014 WIC Facts: Benton County Health Department. Retrieved from [https://public.health.oregon.gov/HealthyPeopleFamilies/wic/Documents/annual/annual\\_benton.pdf](https://public.health.oregon.gov/HealthyPeopleFamilies/wic/Documents/annual/annual_benton.pdf)

- 
- <sup>94</sup> Oregon Health Authority. (2014). 2014 WIC Facts: Lincoln County Health Department. Retrieved from [https://public.health.oregon.gov/HealthyPeopleFamilies/wic/Documents/annual/annual\\_lincoln.pdf](https://public.health.oregon.gov/HealthyPeopleFamilies/wic/Documents/annual/annual_lincoln.pdf)
- <sup>95</sup> Food Share of Lincoln County. (2015). Our Annual Report, Food Share of Lincoln County 2014-15 Highlights. Retrieved from <http://www.foodsharelincolncounty.org/about-us/our-annual-report/>
- <sup>96</sup> U.S. Department of Housing and Urban Development. (2015). Affordable Housing. Retrieved from [http://portal.hud.gov/hudportal/HUD?src=/program\\_offices/comm\\_planning/affordablehousing/](http://portal.hud.gov/hudportal/HUD?src=/program_offices/comm_planning/affordablehousing/)
- <sup>97</sup> Ending Homelessness Advisory Council. (2008). A Home for Hope: A Ten Year Plan to End Homelessness in Oregon. Retrieved from <http://www.oregon.gov/ohcs/pdfs/report-ehac-10-year-action-plan.pdf>
- <sup>98</sup> Health Resources and Services Administration (2015). Uniform Data System Health Center Program Grantee Profiles. Retrieved from <http://bphc.hrsa.gov/uds/datacenter.aspx?q=d&year=2014&state=OR#glist>.
- <sup>99</sup> Virginia Commonwealth University Center on Society and Health. (2014). Health Care: Necessary But Not Sufficient. Retrieved from [http://www.rwjf.org/content/dam/farm/reports/issue\\_briefs/2014/rwjf415715](http://www.rwjf.org/content/dam/farm/reports/issue_briefs/2014/rwjf415715)
- <sup>100</sup> The Henry J. Kaiser Family Foundation. (2012). Focus on Healthcare Disparities: Key Facts. Retrieved from <https://kaiserfamilyfoundation.files.wordpress.com/2012/11/8396-disparities-in-health-and-health-care-five-key-questions-and-answers.pdf>
- <sup>101</sup> The Henry J. Kaiser Family Foundation. (2015). Key Facts about the Uninsured Population. Retrieved from <http://files.kff.org/attachment/fact-sheet-key-facts-about-the-uninsured-population>
- <sup>102</sup> The Henry J. Kaiser Family Foundation. (2013). The Uninsured A Primer 2013-4: How does lack of health insurance affect access to health care? Retrieved from <http://kff.org/report-section/the-uninsured-a-primer-2013-4-how-does-lack-of-insurance-affect-access-to-health-care/>
- <sup>103</sup> Oregon Health Authority, Office of Health Analytics. (2014). Oregon Health Plan: Coordinated Care, Managed Care and Fee for Service Enrollment for January 15, 2014. Retrieved from <http://www.oregon.gov/oha/healthplan/DataReportsDocs/January%202014%20Coordinated%20Care%20Service%20Delivery%20by%20County.pdf>
- <sup>104</sup> Oregon Health Authority, Office of Health Analytics. (2014). State of Oregon: Oregon Health Plan, Medicaid, and CHIP Population by County and Medical Care Delivery System: 15 January 2010. Retrieved from <http://www.oregon.gov/oha/healthplan/DataReportsDocs/January%202010%20Physical%20Health%20Service%20Delivery%20by%20County.pdf>
- <sup>105</sup> Oregon Health Authority, Oregon Health & Science University. (2015). Impacts of the Affordable Care Act on Health Insurance Coverage in Oregon: County Results/Statewide Update. Retrieved from <http://www.ohsu.edu/xd/research/centers-institutes/center-for-health-systems-effectiveness/upload/Health-Insurance-Coverage-in-Oregon-County-Results.pdf>
- <sup>106</sup> Oregon Health Authority, Oregon Health & Science University. (2015). Impacts of the Affordable Care Act on Health Insurance Coverage in Oregon: County Results/Statewide Update. Retrieved from <http://www.ohsu.edu/xd/research/centers-institutes/center-for-health-systems-effectiveness/upload/Health-Insurance-Coverage-in-Oregon-County-Results.pdf>
- <sup>107</sup> Oregon Health Authority, Oregon Health & Science University. (2015). Impacts of the Affordable Care Act on Health Insurance Coverage in Oregon: County Results/Statewide Update. Retrieved from <http://www.ohsu.edu/xd/research/centers-institutes/center-for-health-systems-effectiveness/upload/Health-Insurance-Coverage-in-Oregon-County-Results.pdf>
- <sup>108</sup> U.S. Census Bureau. (2013). Health Insurance Coverage Status, American Community Survey 3-Year estimates, 2011-2013, Table S2701. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_S2701&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_S2701&prodType=table)
- <sup>109</sup> U.S. Census Bureau. (2013). Health Insurance Coverage Status, American Community Survey 3-Year estimates, 2011-2013, Table S2701. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_S2701&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_S2701&prodType=table)
- <sup>110</sup> UCLA Center for Health Policy Research. (2012). Undocumented Immigrants and Health Care Reform. Retrieved from <http://healthpolicy.ucla.edu/publications/Documents/PDF/undocumentedreport-aug2013.pdf>
- <sup>111</sup> U.S. Census Bureau. (2013). Health Insurance Coverage Status, American Community Survey 3-Year estimates, 2011-2013, Table S2701. Retrieved from [http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_13\\_3YR\\_S2701&prodType=table](http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_13_3YR_S2701&prodType=table)
- <sup>112</sup> Alliance for a Justice Society. (2015). Breaking Barriers: Improving Health Insurance Enrollment and Access to Health Care. Retrieved from [http://allianceforajustsociety.org/wp-content/uploads/2015/04/BreakingBarriers\\_Natl\\_sm.pdf](http://allianceforajustsociety.org/wp-content/uploads/2015/04/BreakingBarriers_Natl_sm.pdf)
- <sup>113</sup> Riffkin R. (2014). Cost Still a Barrier Between Americans and Medical Care. *Gallup Poll*. Retrieved from <http://www.gallup.com/poll/179774/cost-barrier-americans-medical-care.aspx>
- <sup>114</sup> Robert Wood Johnson Foundation, County Health Rankings & Roadmaps. (2015). Could not see a doctor due to cost. Retrieved from <http://www.countyhealthrankings.org/app/oregon/2015/measure/additional/87/data>
- <sup>115</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Access to Health Services. Retrieved from <http://www.healthypeople.gov/2020/topics-objectives/topic/Access-to-Health-Services>
- <sup>116</sup> Robert Wood Johnson Foundation. (2012). Health Care Safety Net Resources by State. Retrieved from <http://www.rwjf.org/en/library/research/2012/02/health-care-safety-net-resources-by-state.html>
- <sup>117</sup> Lincoln County Health and Human Services Department. (2015). Lincoln County Health and Human Services Department. Retrieved from <https://public.health.oregon.gov/ProviderPartnerResources/LocalHealthDepartmentResources/Documents/Lincoln.pdf>

- 
- <sup>118</sup> U.S. Department of Health and Human Services, Health Resources and Services Administration. (n.d.). Health Center Data & Reporting. Retrieved from <http://www.bphc.hrsa.gov/datareporting/index.html>
- <sup>119</sup> U.S. Department of Health and Human Services, Health Resources and Services Administration. (n.d.). Health Professional Shortage Areas (HPSAs). Retrieved from <http://bhpr.hrsa.gov/shortage/hpsas/>
- <sup>120</sup> National Association of Community Health Centers, Inc. (n.d.). Migrant and Seasonal Farmworker Access to Health Care Services and Insurance Coverage: Summary Report on Issues, Resources and Potential Solutions. Retrieved from [http://www.cpc.org/cpca/assets/File/Policy-and-Advocacy/Active-Policy-Issues/MSFW/MSFW\\_access\\_to\\_health\\_care\\_services.pdf](http://www.cpc.org/cpca/assets/File/Policy-and-Advocacy/Active-Policy-Issues/MSFW/MSFW_access_to_health_care_services.pdf)
- <sup>121</sup> Oregon Office of Rural Health. (2015). 2015 Areas of Unmet Health Care Need in Rural Oregon Report. Retrieved from <http://www.ohsu.edu/xd/outreach/oregon-rural-health/data/upload/2015-Unmet-Need-Report.pdf>
- <sup>122</sup> Oregon Office of Rural Health. (2015). 2015 Areas of Unmet Health Care Need in Rural Oregon Report. Retrieved from <http://www.ohsu.edu/xd/outreach/oregon-rural-health/data/upload/2015-Unmet-Need-Report.pdf>
- <sup>123</sup> Samaritan Health Services. (2015). Surgical Services. Retrieved from <http://www.samhealth.org/services/surgery/Pages/services.aspx>
- <sup>124</sup> Samaritan Health Services. (2013). Girod Birth Center (SLCH). Retrieved 1 November 2013 from <http://www.samhealth.org/services/womensservices/Pages/girodbirthcenterslch.aspx>
- <sup>125</sup> Samaritan Health Services. (2013). Imagine services by location. Retrieved from [http://www.samhealth.org/SiteCollectionDocuments/imaging-chart\\_05.2011.pdf](http://www.samhealth.org/SiteCollectionDocuments/imaging-chart_05.2011.pdf)
- <sup>126</sup> Gindi, R., Cohen, R., and Kirzinger, W. (2012). Emergency Room Use Among Adults Aged 18–64: Early Release of Estimates From the National Health Interview Survey, January–June 2011. Retrieved from [http://www.cdc.gov/nchs/data/nhis/earlyrelease/emergency\\_room\\_use\\_january-june\\_2011.pdf](http://www.cdc.gov/nchs/data/nhis/earlyrelease/emergency_room_use_january-june_2011.pdf)
- <sup>127</sup> Samaritan Health Services (2015). Hospice and Palliative Care. Retrieved from <http://www.samhealth.org/services/hospicepalliativecare/Pages/default.aspx>
- <sup>128</sup> Oregon Health Authority, Oregon Public Health Division, Oregon Vital Statistics County data 2013. (2013). Table 19. Tobacco-linked deaths by county of residence, Oregon 2013. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/VitalStatistics/annualreports/CountyDataBook/Documents/2013/table19-2013.pdf>
- <sup>129</sup> Macera, C., Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion. (n.d.). Promoting Healthy Eating and Physical Activity for a Healthier Nation. Retrieved from <http://www.cdc.gov/healthyyouth/publications/pdf/pp-ch7.pdf>
- <sup>130</sup> Oregon Health Authority, Oregon Public Health Division, Oregon Vital Statistics County data 2013. (2013). Table 18. Leading causes of death by county of residence, Oregon, 2013. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/VitalStatistics/annualreports/CountyDataBook/Documents/2013/table18-2013.pdf>
- <sup>131</sup> American Cancer Society. (2015). Cancer Facts and Figures 2015. Retrieved from: <http://www.cancer.org/acs/groups/content/@editorial/documents/document/acspc-044552.pdf>
- <sup>132</sup> Oregon Health Authority, Oregon Public Health Division, Tobacco Prevention and Education Program. (2013). Oregon Tobacco Facts: Tobacco-Related Diseases. Retrieved from [https://public.health.oregon.gov/PreventionWellness/TobaccoPrevention/Documents/tobacco\\_facts/tobacco-related\\_diseases.pdf](https://public.health.oregon.gov/PreventionWellness/TobaccoPrevention/Documents/tobacco_facts/tobacco-related_diseases.pdf)
- <sup>133</sup> Oregon Health Authority, Oregon Public Health Division. (2012). Cancer in Oregon. *CD Summary*, 61(19). Retrieved from <http://public.health.oregon.gov/DiseasesConditions/CommunicableDisease/CDSummaryNewsletter/Documents/2012/ohd6119.pdf>
- <sup>134</sup> Oregon Health Authority, Oregon Public Health Division. (2012). Cancer in Oregon. *CD Summary*, 61(19). Retrieved from <http://public.health.oregon.gov/DiseasesConditions/CommunicableDisease/CDSummaryNewsletter/Documents/2012/ohd6119.pdf>
- <sup>135</sup> Oregon Health Authority. (2014). Oregon Tobacco Laws and Policies. Retrieved from <http://public.health.oregon.gov/PreventionWellness/TobaccoPrevention/Documents/tobfacts.pdf>
- <sup>136</sup> American Cancer Society. (2012). Cancer Facts and Figures 2012. Retrieved from: <http://www.cancer.org/acs/groups/content/%40epidemiologysurveillance/documents/document/acspc-031941.pdf>
- <sup>137</sup> Susan G. Komen Foundation. (n.d.). Facts and Statistics. Retrieved from <http://www5.komen.org/BreastCancer/FactsandStatistics.html>
- <sup>138</sup> National Cancer Institute, State Cancer Profiles. (n.d.). Incidence Rates Table, Breast (Female). Retrieved from <http://statecancerprofiles.cancer.gov/incidencerates/index.php?stateFIPS=41&cancer=055&race=00&sex=2&age=001&type=incd#results>
- <sup>139</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Cancer, C-3. Retrieved from <https://www.healthypeople.gov/2020/topics-objectives/topic/cancer/objectives>
- <sup>140</sup> Oregon Health Authority, Oregon State Cancer Registry. (2006). Cancer in Oregon, 2006. Retrieved from <https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Cancer/oscar/Documents/ar2006/ar06.pdf>
- <sup>141</sup> Office of Disease Prevention and Health Promotion, Healthy People 2020 (2014). 2020 Topics & Objectives: Cancer. Retrieved from <http://www.healthypeople.gov/2020/topics-objectives/topic/cancer/objectives>
- <sup>142</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Cancer, C-5. Retrieved from <https://www.healthypeople.gov/2020/topics-objectives/topic/cancer/objectives>

- 
- <sup>143</sup> National Cancer Institute, State Cancer Profiles. (n.d.). Incidence Rates Table, Pancreas. Retrieved from <http://www.statecancerprofiles.cancer.gov/incidencrates/index.php?stateFIPS=41&cancer=040&race=00&sex=0&age=001&type=i ncd#results>
- <sup>144</sup> Oregon Health Authority, Public Health Division, Health Promotion & Chronic Disease Prevention Section. (2013). Diabetes, Heart Disease and Stroke in Oregon. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Documents/OHA8582\\_AllVolumes.pdf](https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Documents/OHA8582_AllVolumes.pdf)
- <sup>145</sup> Franks, P., Winters, P., Tancredi, D., Fiscella, K. (2011). Do changes in traditional coronary heart disease risk factors over time explain the association between socio-economic status and coronary heart disease? *BMC Cardiovascular Disorders*, 11(28). Retrieved from <http://www.biomedcentral.com/1471-2261/11/28>
- <sup>146</sup> Oregon Health Authority, Public Health Division, Office of Disease Prevention and Epidemiology. (2010). Heart Disease and Stroke in Oregon: Update - 2010. Retrieved from [http://public.health.oregon.gov/DiseasesConditions/ChronicDisease/HeartDiseaseStroke/Documents/heartstroke\\_update2010.pdf](http://public.health.oregon.gov/DiseasesConditions/ChronicDisease/HeartDiseaseStroke/Documents/heartstroke_update2010.pdf)
- <sup>147</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Heart Disease Mortality, by Case and County. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>148</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Heart Disease and Stroke, Objectives. Retrieved from <https://www.healthypeople.gov/2020/topics-objectives/topic/heart-disease-and-stroke/objectives>
- <sup>149</sup> American Diabetes Association. (n.d.). Lower Your Risk. Retrieved from <http://www.diabetes.org/are-you-at-risk/lower-your-risk/>
- <sup>150</sup> Oregon Health Authority, Public Health Division, Oregon BRFSS County Combined Dataset. (2008-2011). Table 1: Age-Adjusted and Unadjusted Prevalence of Selected Chronic Conditions among Adults, by County, Oregon 2008-2011. Retrieved from [http://public.health.oregon.gov/BirthDeathCertificates/Surveys/AdultBehaviorRisk/county/Documents/0811/ORCounty\\_conditions.pdf](http://public.health.oregon.gov/BirthDeathCertificates/Surveys/AdultBehaviorRisk/county/Documents/0811/ORCounty_conditions.pdf)
- <sup>151</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Diabetes, Objectives. Retrieved from <http://www.healthypeople.gov/2020/topics-objectives/topic/diabetes/objectives>
- <sup>152</sup> Oregon Health Authority, Public Health Division, Oregon BRFSS County Combined Dataset. (2008-2011). Table 1: Age-Adjusted and Unadjusted Prevalence of Selected Chronic Conditions among Adults, by County, Oregon 2008-2011. Retrieved from [http://public.health.oregon.gov/BirthDeathCertificates/Surveys/AdultBehaviorRisk/county/Documents/0811/ORCounty\\_conditions.pdf](http://public.health.oregon.gov/BirthDeathCertificates/Surveys/AdultBehaviorRisk/county/Documents/0811/ORCounty_conditions.pdf)
- <sup>153</sup> Garland-Forshee, R., Gedman, T., Oregon Health Authority, Public Health Division, Center or Prevention and Health Promotion, Oregon Asthma Program. (2013). The Burden of Asthma in Oregon. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Asthma/Documents/burden/OR\\_Asthma\\_2013.pdf](https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Asthma/Documents/burden/OR_Asthma_2013.pdf)
- <sup>154</sup> Garland-Forshee, R., Gedman, T., Oregon Health Authority, Public Health Division, Center or Prevention and Health Promotion, Oregon Asthma Program. (2013). The Burden of Asthma in Oregon. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Asthma/Documents/burden/OR\\_Asthma\\_2013.pdf](https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Asthma/Documents/burden/OR_Asthma_2013.pdf)
- <sup>155</sup> Garland-Forshee, R., Gedman, T., Oregon Health Authority, Public Health Division, Center or Prevention and Health Promotion, Oregon Asthma Program. (2013). The Burden of Asthma in Oregon. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Asthma/Documents/burden/OR\\_Asthma\\_2013.pdf](https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/Asthma/Documents/burden/OR_Asthma_2013.pdf)
- <sup>156</sup> Oregon Health Authority, Public Health Division, Oregon BRFSS County Combined Dataset. (2008-2011). Table 1: Age-Adjusted and Unadjusted Prevalence of Selected Chronic Conditions among Adults, by County, Oregon 2008-2011. Retrieved from [http://public.health.oregon.gov/BirthDeathCertificates/Surveys/AdultBehaviorRisk/county/Documents/0811/ORCounty\\_conditions.pdf](http://public.health.oregon.gov/BirthDeathCertificates/Surveys/AdultBehaviorRisk/county/Documents/0811/ORCounty_conditions.pdf)
- <sup>157</sup> World Health Organization. (n.d.). Health topics: Infectious diseases. Retrieved from [http://www.who.int/topics/infectious\\_diseases/en/](http://www.who.int/topics/infectious_diseases/en/)
- <sup>158</sup> Oregon Health Authority, Public Health Division. (n.d.). Oregon Disease Reporting: What is Reportable and When. Retrieved from <https://public.health.oregon.gov/DiseasesConditions/CommunicableDisease/ReportingCommunicableDisease/Pages/reportable.aspx>
- <sup>159</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Communicable Disease Reports, Case Count by Disease and County. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>160</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Communicable Disease Reports, Case Count by Disease and County. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>161</sup> Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases, Division of Foodborne, Waterborne, and Environmental Diseases. (2014). Estimates of foodborne illness in the United States. Retrieved from <http://www.cdc.gov/foodborneburden/2011-foodborne-estimates.html>
- <sup>162</sup> Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases, Division of Foodborne, Waterborne, and Environmental Diseases. (2014). Estimates of foodborne illness in the United States. Retrieved from <http://www.cdc.gov/foodborneburden/2011-foodborne-estimates.html>
- <sup>163</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Communicable Disease Reports, Case Count by Disease and County. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>164</sup> Centers for Disease Control and Prevention. (2011). STDs in Women and Infants. Retrieved from <http://www.cdc.gov/std/stats10/womenandinf.htm>



- 
- <sup>165</sup> Oregon Health Authority, Public Health Division. (2014). Chlamydia in Oregon. Retrieved from <https://public.health.oregon.gov/DiseasesConditions/CommunicableDisease/DiseaseSurveillanceData/STD/Documents/9980STD-Chlamydia.pdf>
- <sup>166</sup> Oregon Health Authority, Public Health Division. (2014). Gonorrhea in Oregon. Retrieved from <https://public.health.oregon.gov/DiseasesConditions/CommunicableDisease/DiseaseSurveillanceData/STD/Documents/9987-STD-Gonorrhea.pdf>
- <sup>167</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Communicable Disease Reports, Case Count by Disease and County. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>168</sup> Robert Wood Johnson Foundation, County Health Rankings & Roadmaps. (2013). Lincoln County Snapshot. Retrieved from <http://www.countyhealthrankings.org/app/oregon/2012/Lincoln/county/1/overall/snapshot/by-rank>
- <sup>169</sup> National Archive of Criminal Justice Data. (2012). Uniform Crime Reporting Program Data: County-level detailed arrest and offense data, 2012 (ICPSR 35019). Retrieved from <http://www.icpsr.umich.edu/icpsrweb/NACJD/studies/35019>
- <sup>170</sup> Oregon Health Authority, Public Health Division, Oregon Violent Death Reporting System. (2013). Violent Deaths in Oregon: 2013 Retrieved from <https://public.health.oregon.gov/DiseasesConditions/InjuryFatalityData/Documents/NVDRS/NVDRS-Report.pdf>
- <sup>171</sup> Oregon Health Authority, Public Health Division. (2012). Causes of Death: Leading Causes of Death. Retrieved from <https://public.health.oregon.gov/ProviderPartnerResources/PublicHealthAccreditation/Documents/indicators/leadingcausesofdeath.pdf>
- <sup>172</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Injury and Violence Prevention. Retrieved from <http://www.healthypeople.gov/2020/topics-objectives/topic/injury-and-violence-prevention/national-snapshot?topicId=24>
- <sup>173</sup> Oregon Health Authority, Public Health Division, Injury and Violence Prevention Program. (2013). Injury in Oregon, 2013 Injury Data Report. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/InjuryFatalityData/Documents/Injury\\_in\\_Oregon\\_v2.3.pdf](https://public.health.oregon.gov/DiseasesConditions/InjuryFatalityData/Documents/Injury_in_Oregon_v2.3.pdf)
- <sup>174</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Mental Health. Retrieved from <http://www.healthypeople.gov/2020/leading-health-indicators/2020-lhi-topics/Mental-Health/determinants?tab=determinants>
- <sup>175</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Mental Health. Retrieved from <http://www.healthypeople.gov/2020/leading-health-indicators/2020-lhi-topics/Mental-Health>
- <sup>176</sup> Robert Wood Johnson Foundation, County Health Rankings & Roadmaps. (2014). Lincoln County Snapshot: Poor mental health days. Retrieved from <http://www.countyhealthrankings.org/app/oregon/2014/rankings/Lincoln/county/outcomes/overall/snapshot>
- <sup>177</sup> Oregon Health Authority, Public Health division, Oregon Behavioral Risk Factor Surveillance System. (2013). Oregon Adults in Good Mental Health, Oregon, 2008-2011 (Age-Adjusted). Retrieved from [https://public.health.oregon.gov/BirthDeathCertificates/Surveys/AdultBehaviorRisk/county/Documents/0811/MentalHealthAA\\_0811.pdf](https://public.health.oregon.gov/BirthDeathCertificates/Surveys/AdultBehaviorRisk/county/Documents/0811/MentalHealthAA_0811.pdf)
- <sup>178</sup> Oregon Health Authority, Public Health Division, Oregon Behavioral Risk Factor Surveillance System. (2013). Chronic diseases among Oregon adults, by county, 2010-2013. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Documents/datatables/ORCountyBRFSS\\_diseases.pdf](https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Documents/datatables/ORCountyBRFSS_diseases.pdf)
- <sup>179</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Selected Causes of Death by County. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>180</sup> World Health Organization. (n.d.). The implications for training of embracing: A Life Course Approach to Health. Retrieved from [http://www.who.int/ageing/publications/lifecourse/alc\\_lifecourse\\_training\\_en.pdf](http://www.who.int/ageing/publications/lifecourse/alc_lifecourse_training_en.pdf)
- <sup>181</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Birth Risk Factors: Percent of Births for which Birth Risk Factor is Present. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>182</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Birth Risk Factors: Percent of Births for which Birth Risk Factor is Present. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>183</sup> Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Division of Reproductive Health. (2007). Preventing Smoking and Exposure to Secondhand Smoke Before, During, and After Pregnancy. Retrieved from <http://www.cdc.gov/nccdphp/publications/factsheets/prevention/pdf/smoking.pdf>
- <sup>184</sup> Centers for Disease Control and Prevention, National Center on Birth Defects and Developmental Disabilities, Division of Birth Defects and Developmental Disabilities. (2014). Fetal Alcohol Spectrum Disorders (FASDs). Retrieved from <http://www.cdc.gov/ncbddd/fasd/alcohol-use.html>
- <sup>185</sup> Oregon Health Authority, Public Health Division. (2011). Oregon PRAMS: 2011 Results. Retrieved from <https://public.health.oregon.gov/HealthyPeopleFamilies/DataReports/prams/Documents/OregonPRAMS2011.pdf>
- <sup>186</sup> U.S. Department of Health and Human Services, National Institute of Child Health and Human Development. (2013). What are the factors that put a pregnancy at risk? Retrieved from <http://www.nichd.nih.gov/health/topics/high-risk/conditioninfo/pages/factors.aspx>
- <sup>187</sup> Planned Parenthood Federation of America. (2013). Pregnancy and Childbearing Among U.S. Teens. Retrieved from [http://www.plannedparenthood.org/files/2013/9611/7570/Pregnancy\\_And\\_Childbearing\\_Among\\_US\\_Teens.pdf](http://www.plannedparenthood.org/files/2013/9611/7570/Pregnancy_And_Childbearing_Among_US_Teens.pdf)
- <sup>188</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Birth Risk Factors: Percent of Births for which Birth Risk Factor is Present. Retrieved from <https://ophat.public.health.oregon.gov>

- 
- <sup>189</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Maternal, Infant, and Child Health. Retrieved from <http://www.healthypeople.gov/2020/topics-objectives/topic/maternal-infant-and-child-health/objectives>
- <sup>190</sup> Barker, D.J., Osmond, C. (1986). Infant Mortality, Childhood Nutrition and Ischaemic Heart Disease in England and Wales. *Lancet*, 327(8489), 1077-81. DOI: 10.1111/j.1365-2796.2007.01809.x
- <sup>191</sup> Hack, M., Klein, N., Taylor, H. (1995). Long-term developmental outcomes of low birthweight children. *Future Child*, 5(1), 176-196. DOI: 0.2307/1602514
- <sup>192</sup> World Health Organization. (2015). Preterm birth. Retrieved from <http://www.who.int/mediacentre/factsheets/fs363/en/>
- <sup>193</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Birth Risk Factors: Percent of Births for which Birth Risk Factor is Present. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>194</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Maternal, Infant, and Child Health, Objectives. Retrieved from <http://www.healthypeople.gov/2020/topics-objectives/topic/maternal-infant-and-child-health/objectives>
- <sup>195</sup> American Academy of Pediatrics. (2012). Policy Statement: Breastfeeding and the use of human milk. *Pediatrics*, 129(3), 827-841. Retrieved from <http://pediatrics.aappublications.org/content/129/3/e827>
- <sup>196</sup> Oregon Health Authority, Public Health Division, Oregon WIC Program. (2014). 2014 WIC Facts. Retrieved from [https://public.health.oregon.gov/HealthyPeopleFamilies/wic/Documents/annual/annual\\_all.pdf](https://public.health.oregon.gov/HealthyPeopleFamilies/wic/Documents/annual/annual_all.pdf)
- <sup>197</sup> Centers for Disease Control and Prevention, National Center for Injury Prevention and Control, Division of Violence Prevention. (2014). Adverse Childhood Experiences (ACE) study. Retrieved from <http://www.cdc.gov/ace/outcomes.htm>
- <sup>198</sup> Felitti, V.J., Anda, R.F., Nordenberg, D., Williamson, D.F., Spitz, A.M., Edwards, V., et al. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) study. *American Journal of Preventive Medicine*, 14(4), 245-258. Retrieved from <http://www.theannainstitute.org/ACE%20folder%20for%20website/4RCH.pdf>
- <sup>199</sup> Oregon Health Authority, Addictions and Mental Health Division. (2014). 2014 Oregon Student Wellness Survey: Lincoln County. Retrieved from [https://oregon.pridesurveys.com/dl.php?pdf=Lincoln\\_Co\\_2014.pdf&type=county](https://oregon.pridesurveys.com/dl.php?pdf=Lincoln_Co_2014.pdf&type=county)
- <sup>200</sup> Oregon Health Authority, Addictions and Mental Health Division. (2014). 2014 Oregon Student Wellness Survey: Lincoln County. Retrieved from [https://oregon.pridesurveys.com/dl.php?pdf=Lincoln\\_Co\\_2014.pdf&type=county](https://oregon.pridesurveys.com/dl.php?pdf=Lincoln_Co_2014.pdf&type=county)
- <sup>201</sup> Oregon Health Authority, Addictions and Mental Health Division. (2014). 2014 Oregon Student Wellness Survey: Benton County. Retrieved from [https://oregon.pridesurveys.com/dl.php?pdf=Benton\\_Co\\_2014.pdf&type=county](https://oregon.pridesurveys.com/dl.php?pdf=Benton_Co_2014.pdf&type=county)
- <sup>202</sup> Oregon Department of Human Services, Children, Adults and Families Division. (2015). 2014 Child Welfare Data Book. Retrieved from <http://www.oregon.gov/dhs/children/child-abuse/Documents/2014%20Data%20Book.pdf>
- <sup>203</sup> Oregon Department of Human Services, Children, Adults and Families Division. (2015). 2014 Child Welfare Data Book. Retrieved from <http://www.oregon.gov/dhs/children/child-abuse/Documents/2014%20Data%20Book.pdf>
- <sup>204</sup> Oregon Department of Human Services, Children, Adults and Families Division. (2012). 2011 Child Welfare Data Book. Retrieved from <http://www.oregon.gov/dhs/children/child-abuse/Documents/2011-cw-data-book.pdf>
- <sup>205</sup> Oregon Department of Human Services, Children, Adults and Families Division. (2013). 2012 Foster Care Data Book. Retrieved from <http://www.oregon.gov/dhs/children/child-abuse/Documents/2012-cw-data-book.pdf>
- <sup>206</sup> Oregon Department of Human Services, Children, Adults and Families Division. (2014). 2013 Child Welfare Data Book. Retrieved from <http://www.oregon.gov/dhs/children/child-abuse/Documents/2013%20Data%20Book.pdf>
- <sup>207</sup> Oregon Department of Human Services, Children, Adults and Families Division. (2015). 2014 Child Welfare Data Book. Retrieved from <http://www.oregon.gov/dhs/children/child-abuse/Documents/2014%20Data%20Book.pdf>
- <sup>208</sup> Oregon Department of Health Services. (2015). Foster Care. Retrieved from <http://www.oregon.gov/dhs/children/fostercare/pages/index.aspx>
- <sup>209</sup> Bosworth, K., Espelage, D., Simon, T. (1999). Factors associated with bullying behavior in middle school students. *Journal of Early Adolescence*, 19(3), 341-362. Retrieved from [http://extension.fullerton.edu/professionaldevelopment/assets/pdf/bullying/bullying\\_and\\_middleschool\\_students.pdf](http://extension.fullerton.edu/professionaldevelopment/assets/pdf/bullying/bullying_and_middleschool_students.pdf)
- <sup>210</sup> Oregon Health Authority. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>
- <sup>211</sup> Oregon Health Authority. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>
- <sup>212</sup> World Health Organization. (2014). Mental health: strengthening our response. Retrieved from <http://www.who.int/mediacentre/factsheets/fs220/en/>
- <sup>213</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Suicide Rates, 2009-2013. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>214</sup> Oregon Health Authority, Public Health Division. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>
- <sup>215</sup> Healthy People 2020. (2015). Physical Activity: Objectives. Retrieved from <http://www.healthypeople.gov/2020/topics-objectives/topic/physical-activity/objectives>
- <sup>216</sup> Oregon Health Authority, Public Health Division. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>

- 
- <sup>217</sup> Oregon Health Authority, Public Health Division. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>
- <sup>218</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Physical Activity, PA-8.3.3. Retrieved from <http://www.healthypeople.gov/2020/topics-objectives/topic/physical-activity/objectives>
- <sup>219</sup> U.S. Department of Health & Human Services, National Institutes of Health, National Heart, Lung, and Blood Institute. (2012). What Are the Health Risks of Overweight and Obesity? Retrieved from <http://www.nhlbi.nih.gov/health/health-topics/topics/obe/risks>
- <sup>220</sup> Kennedy, E. (2006). Evidence for Nutritional Benefits in Prolonging Wellness. *The American Journal of Clinical Nutrition*, 83(suppl), 410S-4S. Retrieved from <http://ajcn.nutrition.org/content/83/2/410S.full.pdf>
- <sup>221</sup> Oregon Health Authority, Public Health Division. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>
- <sup>222</sup> National Institute of Dental and Craniofacial Research. (2014). Oral Health in America: A Report of the Surgeon General (Executive Summary). Retrieved from <http://www.nidcr.nih.gov/DataStatistics/SurgeonGeneral/Report/ExecutiveSummary.htm>
- <sup>223</sup> Oregon Public Health Authority, Public Health Division. (n.d.). Oral Health for Infants and Children. Retrieved from <http://public.health.oregon.gov/PreventionWellness/oralhealth/Pages/child.aspx>
- <sup>224</sup> Oregon Health Authority, Public Health Division. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>
- <sup>225</sup> Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Division of Oral Health. (2008). Populations Receiving Optimally Fluoridated Public Drinking Water – United States, 1992-2006. *MMWR*, 57(27), 737-741. Retrieved from <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5727a1.htm>
- <sup>226</sup> Oregon Health & Science University, Portland State University. (2013). State of Our Health 2013: Key Health Indicators for Oregonians. Retrieved from <https://www.ohsu.edu/xd/education/student-services/about-us/provost/upload/State-of-Our-Health-2013-monograph.pdf>
- <sup>227</sup> Benton, Lincoln, Lincoln Regional Oral Health Coalition. (2015). Oral Health Needs in Benton, Lincoln and Lincoln Counties: An Assessment. Retrieved from <http://www.samhealth.org/SiteCollectionDocuments/CommunitySupport/Regional%20NeedsAssessment2015.pdf>
- <sup>228</sup> National Council on Alcoholism and Drug Dependence, Inc. (2015). Facts About Drugs. Retrieved from <https://ncadd.org/about-addiction/faq/facts-about-drugs>
- <sup>229</sup> National Institutes of on Alcohol Abuse and Alcoholism. (2006). No. 68: Young Adult Drinking [factsheet]. Retrieved from <http://pubs.niaaa.nih.gov/publications/aa68/aa68.htm>
- <sup>230</sup> Oregon Health Authority, Public Health Division. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>
- <sup>231</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Substance Abuse. Retrieved from <https://www.healthypeople.gov/2020/topics-objectives/topic/substance-abuse/objectives>
- <sup>232</sup> U.S. Department of Health and Human Services. (2012). Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General. Retrieved from <http://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-report.pdf>
- <sup>233</sup> Oregon Health Authority, Public Health Division. (2007). Oregon Healthy Teens Survey: 2007 Results, County Level Results. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/2007/Pages/sda.aspx>
- <sup>234</sup> Oregon Health Authority, Public Health Division. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>
- <sup>235</sup> Oregon Health Authority, Public Health Division. (2015). Oregon Healthy Teens Survey: 2014/2015 School Year Results, 2015 County Reports. Retrieved from <https://public.health.oregon.gov/BirthDeathCertificates/Surveys/OregonHealthyTeens/Pages/2015.aspx>
- <sup>236</sup> Oregon Department of Human Services: Child Welfare Program. (2015). Striving to meet the need: Summary of services provided by sexual and domestic violence programs in Oregon. Retrieved from <http://www.oregon.gov/dhs/abuse/domestic/Documents/2014-dv-annual-report.pdf>
- <sup>237</sup> My Sister's Place. (2015). Retrieved from <http://www.mysistersplace.us/>
- <sup>238</sup> My Sister's Place. (2015). Retrieved from <http://www.mysistersplace.us/>
- <sup>239</sup> The National Child Traumatic Stress Network. (2015). Children and Domestic Violence. Retrieved from <http://www.nctsn.org/content/children-and-domestic-violence>
- <sup>240</sup> U.S. Department of Health and Human Services, Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. (2010). Results from the 2010 National Survey on Drug Use and Health: Mental Health Findings. Retrieved from <http://www.samhsa.gov/data/sites/default/files/NSDUHmhfr2010/NSDUHmhfr2010.htm#fig3-1>
- <sup>241</sup> Oregon Health Authority, Office of Health Analytics and Addictions and Mental Health Division. (2012). Lincoln County's Epidemiological Data on Alcohol, Drugs and Mental Health 2000 to 2012. Retrieved from <http://www.oregon.gov/oha/amh/sew/CountyReports/Lincoln%20County%20-%20Epidemiological%20Data%20on%20Alcohol,%20Drugs%20and%20Mental%20Health%202000%20to%202012.pdf>
- <sup>242</sup> Oregon Health Authority, Office of Health Analytics and Addictions and Mental Health Division. (2012). Benton County's Epidemiological Data on Alcohol, Drugs and Mental Health 2000 to 2012. Retrieved from <http://www.oregon.gov/oha/amh/sew/CountyReports/Benton%20County%20-%20Epidemiological%20Data%20on%20Alcohol,%20Drugs%20and%20Mental%20Health%202000%20to%202012.pdf>
- <sup>243</sup> Oregon Health Authority, Office of Health Analytics and Addictions and Mental Health Division. (2012). Lincoln County's Epidemiological Data on Alcohol, Drugs and Mental Health 2000 to 2012. Retrieved from

- 
- <http://www.oregon.gov/oha/amh/sew/CountyReports/Lincoln%20County%20-%20Epidemiological%20Data%20on%20Alcohol,%20Drugs%20and%20Mental%20Health%202000%20to%202012.pdf>
- <sup>244</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Suicide rates, 2009-2013. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>245</sup> Oregon Health Authority, Public Health Division, Oregon Violent Death Reporting System, Injury and Violence Prevention Program, Center for Prevention and Health Promotion. (2012). Suicides in Oregon: Trends and Risk Factors 2012 Report. Retrieved from <http://public.health.oregon.gov/DiseasesConditions/InjuryFatalityData/Documents/NVDRS/Suicide%20in%20Oregon%202012%20Report.pdf>
- <sup>246</sup> Oregon Health Authority, Oregon Public Health Assessment Tool. (2015). Suicide rates, 2009-2013. Retrieved from <https://ophat.public.health.oregon.gov>
- <sup>247</sup> Centers for Disease Control and Prevention, Division of Nutrition, Physical Activity, and Obesity. (2015). Physical Activity Basics. Retrieved from <http://www.cdc.gov/physicalactivity/basics/index.htm>
- <sup>248</sup> Oregon Health Authority, Public Health Division, Health Promotion and Chronic Disease Prevention Program. (2012). Oregon Overweight, Obesity, Physical Activity and Nutrition Facts. Retrieved from [https://public.health.oregon.gov/PreventionWellness/PhysicalActivity/Documents/Oregon\\_PANfactst\\_2012.pdf](https://public.health.oregon.gov/PreventionWellness/PhysicalActivity/Documents/Oregon_PANfactst_2012.pdf)
- <sup>249</sup> Oregon Health Authority, Public Health Division, Health Promotion and Chronic Disease Prevention. (2015). Health risk and protective factors among Oregon adults, by county, 2010-2013. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Documents/datatables/ORCountyBRFSS\\_riskfactors.pdf](https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Documents/datatables/ORCountyBRFSS_riskfactors.pdf)
- <sup>250</sup> Oregon Health Authority, Public Health Division, Health Promotion and Chronic Disease Prevention Program. (2012). Oregon Overweight, Obesity, Physical Activity and Nutrition Facts. Retrieved from [https://public.health.oregon.gov/PreventionWellness/PhysicalActivity/Documents/Oregon\\_PANfactst\\_2012.pdf](https://public.health.oregon.gov/PreventionWellness/PhysicalActivity/Documents/Oregon_PANfactst_2012.pdf)
- <sup>251</sup> Oregon Health Authority, Public Health Division, Health Promotion and Chronic Disease Prevention. (2015). Health risk and protective factors among Oregon adults, by county, 2010-2013. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Documents/datatables/ORCountyBRFSS\\_riskfactors.pdf](https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Documents/datatables/ORCountyBRFSS_riskfactors.pdf)
- <sup>252</sup> Oregon Health Authority, Public Health Division. (2014). Oregon state health profiles, Overweight and obesity among adults and youth. Retrieved from <http://public.health.oregon.gov/About/Pages/HealthStatusIndicators.aspx>
- <sup>253</sup> National Institute on Drug Abuse. (2014). Adolescents and Young Adults. Retrieved from <http://www.drugabuse.gov/publications/research-reports/prescription-drugs/trends-in-prescription-drug-abuse/adolescents-young-adults>
- <sup>254</sup> Centers for Disease Control and Prevention, National Center for Injury Prevention and Control, Division of Unintentional Injury Prevention. (2015). Injury Prevention and Control: Prescription Drug Overdose, Risk Factors for Prescription Painkiller Abuse and Overdose. Retrieved from <http://www.cdc.gov/drugoverdose/epidemic/riskfactors.html>
- <sup>255</sup> Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Division of Adult and Community Health. (2004). Sociodemographic Differences in Binge Drinking Among Adults. *MMWR*, 53(12); 301-304. Retrieved from <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5312a1.htm>
- <sup>256</sup> National Institutes on Alcohol Abuse and Alcoholism. (n.d.). Overview of Alcohol Consumption. Retrieved from <http://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption>
- <sup>257</sup> U.S. Department of Health and Human Services, National Institute of Health, National Institute of Alcohol Abuse and Alcoholism. (2004). No. 3, NIAAA council approves definition of binge drinking. Retrieved from [http://pubs.niaaa.nih.gov/publications/Newsletter/winter2004/Newsletter\\_Number3.pdf](http://pubs.niaaa.nih.gov/publications/Newsletter/winter2004/Newsletter_Number3.pdf)
- <sup>258</sup> Oregon Health Authority, Public Health Division. (2013). Health risk and protective factors among Oregon adults, by county, 2010-2013. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Documents/datatables/ORCountyBRFSS\\_riskfactors.pdf](https://public.health.oregon.gov/DiseasesConditions/ChronicDisease/DataReports/Documents/datatables/ORCountyBRFSS_riskfactors.pdf)
- <sup>259</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Substance Abuse, Objectives. Retrieved from <https://www.healthypeople.gov/2020/topics-objectives/topic/substance-abuse/objectives>
- <sup>260</sup> U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. (2015). Healthy People 2020: Tobacco, Overview & Impact. Retrieved from <https://www.healthypeople.gov/2020/leading-health-indicators/2020-1hi-topics/Tobacco>
- <sup>261</sup> Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. (2014). Health Effects of Secondhand Smoke. Retrieved from [http://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/secondhand\\_smoke/health\\_effects/index.htm](http://www.cdc.gov/tobacco/data_statistics/fact_sheets/secondhand_smoke/health_effects/index.htm)
- <sup>262</sup> Oregon Health Authority, Public Health Division, Injury and Violence Prevention Program. (2015). Falls among older adults in Oregon. Retrieved from [https://public.health.oregon.gov/DiseasesConditions/InjuryFatalityData/Documents/Fact%20Sheets/Falls\\_Older\\_Adults\\_2015v0226\\_2015.pdf](https://public.health.oregon.gov/DiseasesConditions/InjuryFatalityData/Documents/Fact%20Sheets/Falls_Older_Adults_2015v0226_2015.pdf)
- <sup>263</sup> Oregon Health Authority, Public Health Division, Injury and Violence Prevention Program. (2015). Falls among older adults in Oregon. Retrieved from

---

[https://public.health.oregon.gov/DiseasesConditions/InjuryFatalityData/Documents/Fact%20Sheets/Falls\\_Older\\_Adults\\_2015v02262015.pdf](https://public.health.oregon.gov/DiseasesConditions/InjuryFatalityData/Documents/Fact%20Sheets/Falls_Older_Adults_2015v02262015.pdf)

<sup>264</sup> County Health Rankings. (2016). Retrieved from

<http://www.countyhealthrankings.org/app/oregon/2016/rankings/Lincoln/county/outcomes/overall/snapshot>

<sup>265</sup> County Health Rankings. (2016). Retrieved from

<http://www.countyhealthrankings.org/app/oregon/2016/rankings/Lincoln/county/outcomes/overall/snapshot>